

Lattice Point Asymptotics and Volume Growth on Teichmüller space.

Jayadev Athreya*, Alexander Bufetov†, Alex Eskin‡ and Maryam Mirzakhani§

February 28, 2010

1 Introduction.

In this note, we apply some of the ideas of the Ph.D. Thesis of G. A. Margulis [Mar70] to Teichmüller space. Let $g \geq 2$, let $\mathcal{M}_g$ and $\mathcal{T}_g$ denote the moduli space and the Teichmüller space of compact Riemann surfaces of genus g . Let $\mathcal{Q}^1\mathcal{M}_g$ (resp. $\mathcal{Q}^1\mathcal{T}_g$) denote the bundle of unit-area quadratic differentials on $\mathcal{M}_g$ (resp. $\mathcal{T}_g$). Let Γ be the mapping class group, so e.g. $\mathcal{Q}^1\mathcal{M}_g = \mathcal{Q}^1\mathcal{T}_g/\Gamma$.

The space $\mathcal{Q}^1\mathcal{T}_g$ carries a natural normalized smooth measure μ , preserved by the action of Γ and such that $\mu(\mathcal{Q}^1\mathcal{M}_g) = 1$ ([Mas82], [Ve82]). We have a natural projection

$$\pi : \mathcal{Q}^1\mathcal{T}_g \rightarrow \mathcal{T}_g$$

and we set $\mathbf{m} = \pi_*\mu$.

Let $x, y \in \mathcal{T}_g$ be arbitrary points, and let $B_R(x)$ be the ball of radius R in the Teichmüller space in the Teichmüller metric, centered at the point x .

Our main results are the following two theorems:

Theorem 1.1 (Lattice Point Asymptotics) *As $R \rightarrow \infty$,*

$$|\Gamma y \cap B_R(x)| \sim \frac{1}{h} \Lambda(x) \Lambda(y) e^{hR},$$

where $h = 6g - 6$ is the entropy of the Teichmüller geodesic flow, and Λ is the Hubbard-Masur function defined in §3. (Here and below, the notation $A \sim B$ means that $A/B \rightarrow 1$ as $R \rightarrow \infty$).

Theorem 1.2 (Volume Asymptotics) *As $R \rightarrow \infty$,*

$$\mathbf{m}(B_R(x)) \sim \frac{1}{h} e^{hR} \Lambda(x) \cdot \int_{\mathcal{M}_g} \Lambda(y) d\mathbf{m}(y).$$

*partially supported by NSF grants DMS 0603636 and DMS 0244542.

†partially supported by NSF grant DMS 0604386, by the Edgar Odell Lovett Fund at Rice University and by the Programme on Mathematical Control Theory of the Presidium of the Russian Academy of Sciences.

‡partially supported by NSF grants DMS 0244542 and DMS 0604251.

§partially supported by the Clay foundation and by NSF grant DMS 0804136.

We also prove versions of the above theorems in “sectors”, see Theorem 7.1, Theorem 7.2 and Theorem 10.2 below.

Acknowledgments. The authors are grateful to Giovanni Forni, Vadim Kaimanovich, Yair Minsky and Kasra Rafi for useful discussions, and especially to Howard Masur for his help with all parts of the paper, in particular the appendix. We are also grateful to the Institute for Advanced Study, IHES and MSRI for their support.

2 The Margulis property of uniform expansion of conditional measures.

Let g_t be the Teichmüller flow on $\mathcal{Q}^1\mathcal{T}_g$. Note that g_t commutes with the action of Γ and preserves the measure μ . Recall that a quadratic differential q is uniquely determined by its imaginary and real measured foliations $\eta^-(q)$ given by $\text{Im}(q^{1/2})$ and $\eta^+(q)$ given by $\text{Re}(q^{1/2})$.

The flow g_t admits the following foliations:

1. $\mathcal{F}^{ss}$, whose leaves are sets of the form $\{q : \eta^+(q) = \text{const}\}$;
2. $\mathcal{F}^{uu}$, whose leaves are sets of the form $\{q : \eta^-(q) = \text{const}\}$.

In other words, for $q_0 \in \mathcal{Q}^1\mathcal{T}_g$, a leaf of $\mathcal{F}^{ss}$ is given by

$$\alpha^{ss}(q_0) = \{q \in \mathcal{Q}^1\mathcal{T}_g : \eta^+(q) = \eta^+(q_0)\},$$

and a leaf of $\mathcal{F}^{uu}$ is given by

$$\alpha^{uu}(q_0) = \{q \in \mathcal{Q}^1\mathcal{T}_g : \eta^-(q) = \eta^-(q_0)\}.$$

Note that the foliations $\mathcal{F}^{ss}$, $\mathcal{F}^{uu}$ are invariant under both g_t and Γ ; in particular, they descend to the moduli space $\mathcal{Q}^1\mathcal{M}_g$.

We also consider the foliation $\mathcal{F}^u$ whose leaves are defined by

$$\alpha^u(q) = \bigcup_{t \in \mathbb{R}} g_t \alpha^{uu}(q)$$

and $\mathcal{F}^s$ whose leaves are defined by

$$\alpha^s(q) = \bigcup_{t \in \mathbb{R}} g_t \alpha^{ss}(q).$$

Denote by $p : \mathcal{MF} \rightarrow \mathcal{PMF}$ the natural projection from the space of measured foliations onto the space of projective measured foliations. Now we may write

$$\alpha^s(q_0) = \{q \in \mathcal{Q}^1\mathcal{T}_g : p(\eta^+(q)) = p(\eta^+(q_0))\};$$

$$\alpha^u(q_0) = \{q \in \mathcal{Q}^1\mathcal{T}_g : p(\eta^-(q)) = p(\eta^-(q_0))\}.$$

The foliations $\mathcal{F}^u$ and $\mathcal{F}^{ss}$ form a complementary pair in the sense of Margulis [Mar70] (so do $\mathcal{F}^s$ and $\mathcal{F}^{uu}$, but the first pair will be more convenient for us). Note that the foliations $\mathcal{F}^{ss}$, $\mathcal{F}^s$, $\mathcal{F}^u$, $\mathcal{F}^{uu}$

are, respectively, the strongly stable, stable, unstable, strongly unstable foliations for the Teichmüller flow in the sense of Veech [Ve86] and Forni [Fo02]. (See Theorem 8.12 below for further results in this direction).

Conditional measures. The main observation, lying at the center of our construction, is the following. Each leaf α^+ of the foliation $\mathcal{F}^{uu}$, as well as each leaf α^- of the foliation $\mathcal{F}^{ss}$ carries a globally defined conditional measure μ_{α^+} , invariant under the action the mapping class group, and having, moreover, the following property:

1. $(g_t)_*\mu_{\alpha^+} = \exp(ht)\mu_{g_t\alpha^+}$;
2. $(g_t)_*\mu_{\alpha^-} = \exp(-ht)\mu_{g_t\alpha^-}$.

where $h = 6g - 6$ is the entropy of the flow g_t on $\mathcal{Q}^1\mathcal{M}_g$ with respect to the smooth measure μ .

The measures $\mu_{\alpha^\pm}$ may be constructed as follows. Let ν denote the Thurston measure on $\mathcal{MF}$. Note that each leaf of $\mathcal{F}^s$ is homeomorphic to an open subset of $\mathcal{MF}$ via the map η^- . We can thus define the conditional measure on $\mathcal{F}^s$ to be the pullback of ν . Similarly one can define the conditional measure on $\mathcal{F}^u$. Now to define the measures $\mu_{\alpha^\pm}$ we must restrict the conditional measure from a leaf of $\mathcal{F}^s$ to a leaf of $\mathcal{F}^{ss}$ (and also from a leaf of $\mathcal{F}^u$ to a leaf of $\mathcal{F}^{uu}$). This can be done explicitly in the following way: for a subset $E \subset \mathcal{MF}$, let $Cone(E)$ denote the cone based at the origin and ending at E (i.e. the union of all the line segments connecting the origin and points of E). We write $\bar{\nu}(E)$ to denote $\nu(Cone(E))$. Now for a set $F \subset \alpha^{ss}(q_0)$, we define $\mu_{\alpha^-}(F) = \bar{\nu}(\eta^-(F))$. Similarly, for $F \subset \alpha^{uu}(q_0)$, $\mu_{\alpha^+}(F)$ is defined to be $\bar{\nu}(\eta^+(F))$.

In particular, if α_1 and α_2 are two leaves of the foliation $\mathcal{F}^u$ and $U_1 \subset \alpha_1$ and $U_2 \subset \alpha_2$ are chosen in such a way that $\eta^+(U_1) = \eta^+(U_2)$, then we have $\mu_{\alpha_1}(U_1) = \mu_{\alpha_2}(U_2)$. The equality $\eta^+(U_1) = \eta^+(U_2)$ is equivalent to the statement that U_1 may be taken to U_2 by holonomy along the leaves of the strongly stable foliation $\mathcal{F}^{ss}$; the equality $\mu_{\alpha_1}(U_1) = \mu_{\alpha_2}(U_2)$ thus means that the smooth measure μ has the property of *holonomy invariance* with respect to the pair of foliations $(\mathcal{F}^u, \mathcal{F}^{ss})$.

This construction allows us to apply the arguments of G.A. Margulis and to compute the asymptotics for the volume of a ball of growing radius in Teichmüller space as well as the asymptotics of the number of elements in the intersection of a ball with the orbit of the mapping class group. The approach is similar to that of [EMc93].

3 The Hubbard-Masur function.

Extremal lengths. Let x be a Riemann surface. Then the extremal length of a simple closed curve γ on x is defined by

$$\text{Ext}_\gamma(x) = \sup_\rho \frac{\ell_\gamma(\rho)^2}{\text{Area}(x, \rho)},$$

where the supremum is taken over all metrics ρ conformally equivalent to x , and $\ell_\gamma(\rho)$ denotes the length of γ in the metric ρ . The extremal length can be extended continuously from the space of simple closed curves to the space of $\mathcal{MF}$ of measured foliations, in such a way that $\text{Ext}_{t\beta}(x) = t^2\text{Ext}_\beta(x)$.

The Hubbard-Masur Theorem. The Hubbard-Masur Theorem [HuMas79] states that given any point $x \in \mathcal{T}_g$ and any measured foliation $\beta \in \mathcal{MF}$, there exists a unique holomorphic quadratic differential q on x such that $\eta^+(q) = \beta$. We also have the identity $\text{Area}(q) = \text{Ext}_\beta(x)$.

The measure s_x and the multiple zero locus. For $x \in \mathcal{T}_g$, we consider the unit (co)-tangent sphere $S(x) = \{q \in \mathcal{Q}^1\mathcal{T}_g : \pi(q) = x\}$ at the point x . The conditional measure of μ on the sphere $S(x)$ will be denoted by s_x .

Let $\mathcal{P}(1, \dots, 1) \subset \mathcal{Q}^1\mathcal{T}_g$ denote the subset where the all zeroes of the quadratic differential are distinct. $\mathcal{P}(1, \dots, 1)$ is called the *principal stratum*. Its complement in $\mathcal{Q}^1\mathcal{T}_g$ is the *multiple zero locus*. We will need the following:

Theorem 3.1 *For any $x \in \mathcal{T}_g$, the measure s_x gives zero weight to the multiple zero locus.*

Proof. See appendix A.

The Hubbard-Masur function. Let $q \in \mathcal{Q}^1\mathcal{T}_g$ and let $\alpha^u(q)$ be the leaf of the foliation $\mathcal{F}^u$. By the Hubbard-Masur Theorem, the projection π induces a continuous bijection between $\alpha^u(q)$ and $\mathcal{T}_g$ which is smooth away from the multiple zero locus. The mapping π thus takes the globally defined conditional measure $\mu_{\alpha^u(q)}$ on the fiber $\alpha^u(q)$ to a measure on the Teichmüller space; the resulting measure $\pi_*(\mu_{\alpha^u(q)})$ is absolutely continuous with respect to the smooth measure $\mathbf{m}$. Furthermore, by Theorem 3.1, the measure $\mathbf{m}$ is also absolutely continuous with respect to $\pi_*(\mu_{\alpha^u(q)})$; indeed, away from the multiple zero locus the mapping π is smooth with a smooth inverse, and the multiple zero locus has measure 0 by Theorem 3.1.

We may therefore consider the corresponding Radon-Nikodym derivative. Introduce a function $\lambda^+ : \mathcal{Q}_g \rightarrow \mathbb{R}$ by the formula

$$\lambda^+(q) = \frac{d\mathbf{m}}{d(\pi_*(\mu_{\alpha^u(q)}))}.$$

Similarly, we define $\lambda^- : \mathcal{Q}_g \rightarrow \mathbb{R}$ via the formula

$$\lambda^-(q) = \frac{d\mathbf{m}}{d(\pi_*(\mu_{\alpha^s(q)}))}.$$

(In the above definitions, the Radon-Nikodym derivatives are evaluated at $\pi(q)$).

We set

$$\Lambda(x) = \int_{S(x)} \lambda^+(q) ds_x(q) = \int_{S(x)} \lambda^-(q) ds_x(q).$$

Note that the functions $\lambda^+, \lambda^-, \Lambda$ are Γ -invariant. We call λ_+, λ_- and Λ the *Hubbard-Masur function*.

Note that by the Hubbard-Masur theorem, η^- (or η^+) defines a homeomorphism the space of all quadratic differentials at x (with arbitrary area) and the space $\mathcal{MF}$. This homeomorphism restricts to a homeomorphism between $S(x)$ and the set

$$E_x = \{\beta \in \mathcal{MF} : \text{Ext}_\beta(x) = 1\}$$

where $\text{Ext}_\beta(x)$ is the extremal length at x of the measured foliation β . Let $\delta_x^\pm : E_x \rightarrow S(x)$ denote the inverse of $\eta^\pm$.

It is easy to see that the functions $\lambda_\pm$ are smooth on the complement of the multiple zero locus.

Proposition 3.2 (Properties of $\lambda^+, \lambda^-, \Lambda$) .

- (i) $\lambda^+(q) = \frac{d(\delta_x^-)_*(\bar{\nu})}{ds_x}(q)$, and $\lambda^-(q) = \frac{d(\delta_x^+)_*(\bar{\nu})}{ds_x}(q)$.
- (ii) $\Lambda(x) = \bar{\nu}(\eta^-(S(x))) = \bar{\nu}(\eta^+(S(x)))$.
- (iii) $\Lambda(x) = \bar{\nu}(E_x) = \nu(\{\beta \in \mathcal{MF} : \text{Ext}_\beta(x) \leq 1\})$.
- (iv) $\Lambda(x) = \nu(\{\beta \in \mathcal{ML} : \text{Ext}_\beta(x) \leq 1\})$, where $\mathcal{ML}$ is the space of measured laminations.

Proof. The property (i) follows from the fact that $d\mu = d\alpha^u d\alpha^{ss}$, the fact that $\eta_*^-(d\alpha^{ss}) = \bar{\nu}$, and the fact that if we write for $q \in \mathcal{Q}^1\mathcal{T}_g$, $q = (x, v)$ where $x = \pi(q)$ and $v \in S(x)$ then $d\mu(q) = d\mathbf{m}(x)d_{s_x}(v)$. The property (ii) follows from (i) after making the change of variable $q \rightarrow \eta^-(q)$ in the definition of Λ . The property (iii) follows from the fact that the image $\eta^-(S(x))$ consists exactly of E_x . Finally (iv) is easily seen to be equivalent to (iii). $\square$

The following is proved in §10:

Theorem 3.3 (Boundedness of Λ) *There exists a constant M such that $\Lambda(x) \leq M$ for all $x \in \mathcal{T}_g$.*

For another interpretation of Λ in terms of asymptotics of the number of multicurves, see (51) below.

4 The Hodge Norm

4.1 Definition and Basic Properties

Let X be a Riemann surface. By definition, X has a complex structure. Let $\mathcal{H}_X$ denote the set of holomorphic 1-forms on X . One can define *Hodge inner product* on $\mathcal{H}_X$ by

$$\langle \omega, \eta \rangle = \int_X \omega \wedge \bar{\eta}.$$

We have a natural map $r : H^1(X, \mathbb{R}) \rightarrow \mathcal{H}_X$ which sends a cohomology class $\lambda \in H^1(X, \mathbb{R})$ to the holomorphic 1-form $r(\lambda) \in \mathcal{H}_X$ such that the real part of $r(\lambda)$ (which is a harmonic 1-form) represents λ . We can thus define the Hodge inner product on $H_1(X, \mathbb{R})$ by $\langle \lambda_1, \lambda_2 \rangle = \langle r(\lambda_1), r(\lambda_2) \rangle$. We have

$$\langle \lambda_1, \lambda_2 \rangle = \int_X \lambda_1 \wedge * \lambda_2,$$

where $*$ denotes the Hodge star operator. We denote the associated norm by $\|\cdot\|$. This is the *Hodge norm*.

Let α be a homology class in $H_1(X, \mathbb{R})$. We can define the cohomology class $*c_\alpha \in H^1(X, \mathbb{R})$ so that for all homology classes $\beta \in H_1(X, \mathbb{R})$,

$$\int_\beta *c_\alpha = I(\beta, \alpha),$$

where $I(\cdot, \cdot)$ denotes algebraic intersection number. Then for any $\alpha, \beta \in H_1(X, \mathbb{R})$,

$$\langle c_\alpha, *c_\beta \rangle = \int_X c_\alpha \wedge c_\beta = \int_X *c_\alpha \wedge *c_\beta = I(\alpha, \beta) = \int_\alpha *c_\beta$$

Since elements of the form $*c_\beta$ span $H^1(X, \mathbb{R})$, it follows that for any $\omega \in H^1(X, \mathbb{R})$,

$$\langle c_\alpha, \omega \rangle = \int_\alpha \omega.$$

We note that $*c_\alpha$ is a purely topological construction which depends only on α , but c_α depends also on the complex structure of X .

4.2 The Hodge Norm and the Hyperbolic Metric.

Let X be a Riemann surface. For a simple closed curve α on X , we let $\ell_\sigma(\alpha)$ denote the length of the geodesic representative of α in the hyperbolic metric which is in the conformal class of X .

Fix $\epsilon_0 > 0$ be such that any two curves of hyperbolic length less than ϵ_0 must be disjoint.

Theorem 4.1 *For any constant $D > 1$ there exists a constant $c > 1$ such that for any simple closed curve α with $\ell_\sigma(\alpha) < D$,*

$$\frac{1}{c} \ell_\sigma(\alpha)^{1/2} \leq \|c_\alpha\| < c \ell_\sigma(\alpha)^{1/2}. \quad (1)$$

Furthermore, if $\ell_\sigma(\alpha) < \epsilon_0$ and β is the shortest simple closed curve crossing α , then

$$\frac{1}{c} \ell_\sigma(\alpha)^{-1/2} \leq \|c_\beta\| < c \ell_\sigma(\alpha)^{-1/2}.$$

Proof. Let $\alpha_1, \dots, \alpha_k$ be the curves with hyperbolic length less than ϵ_0 . For $1 \leq i \leq k$, let β_i be the shortest curve with $i(\alpha_i, \beta_i) = 1$, where $i(\cdot, \cdot)$ denotes the geometric intersection number. It is enough to prove (1) for the α_i and the β_i (the estimate for other moderate length curves follows from a compactness argument).

We can find a collar region around α_i as follows: take two annuli $\{z_i : 1 > |z_i| > |t_i|^{1/2}\}$ and $\{w_i : 1 > w_i > |t_i|^{1/2}\}$ and identify the inner boundaries via the map $w_i = t_i/z_i$. (This coordinate system is used in e.g. [Fa73, Chapter 3], also [Mas76], [Fo02], and elsewhere). The hyperbolic metric σ in the collar region is approximately $|dz|/(|z| \log |z|)$. Then $\ell_\sigma(\alpha_i) \approx 1/|\log t_i|$. By [Fa73, Chapter 3] any holomorphic 1-form ω can be written in the collar region as

$$a_0(z_i + t_i/z_i, t_i) + \frac{a_1(z_i + t_i/z_i, t_i)}{z_i},$$

where a_0 and a_1 are holomorphic. This implies that as $t_i \rightarrow 0$,

$$\omega = \frac{a}{z_i} + h(z_i) + O(t_i/z_i^2)$$

where h is a holomorphic function which remains bounded as $t_i \rightarrow 0$, and the implied constant is bounded as $t_i \rightarrow 0$. (Note that when $|z_i| \geq |t_i|^{1/2}$, $|t_i/z_i^2| \leq 1$). Now from the condition $\int_{\alpha_i} *c_{\beta_i} = 1$ we see that on the collar of α_j ,

$$c_{\beta_i} + i *c_{\beta_i} = \frac{\delta_{ij}}{(2\pi)z_j} + h_{ij}(z_j) + O(t_j/z_j^2), \quad (2)$$

where the h_{ij} are holomorphic and bounded as $t_j \rightarrow 0$. (We use the notation $\delta_{ij} = 1$ if $i = j$ and zero otherwise). Also from the condition $\int_{\beta_i} *c_{\alpha_i} = 1$ we have

$$c_{\alpha_i} + i *c_{\alpha_i} = \frac{1}{|\log t_j|} \left(\frac{\delta_{ij}}{z_j} + s_{ij}(z_j) + O(t_j/z_i^2) \right), \quad (3)$$

where s_{ij} also remains holomorphic and is bounded as $t_j \rightarrow 0$. Then,

$$\|c_{\beta_i}\|^2 = \|*c_{\beta_i}\|^2 = \int_X c_{\beta_i} \wedge *c_{\beta_i} \approx \int_0^{2\pi} \int_{t_i}^1 \frac{1}{4\pi^2 r^2} r dr d\theta \approx \frac{|\log t_i|}{4\pi}$$

and

$$\|c_{\alpha_i}\|^2 = \|*c_{\alpha_i}\|^2 = \int_X c_{\alpha_i} \wedge *c_{\alpha_i} \approx \int_0^{2\pi} \int_{t_i}^1 \frac{1}{(\log t_i)^2 r^2} r dr d\theta \approx \frac{2\pi}{|\log t_i|}$$

□

4.3 The Hodge Norm and Extremal Length.

We recall the following theorem ([Ac60], [Bl61]):

Theorem 4.2 (Accola-Blatter) *For any Riemann surface X and any $\lambda \in H_1(X, \mathbb{R})$,*

$$\|c_\lambda\|^2 = \sup_\rho \inf_{\gamma \in [\lambda]} \frac{\ell_\rho(\gamma)^2}{\text{Area}(\rho)},$$

where the supremum is over the metrics ρ which are in the conformal class of X , and the infimum is over all the multicurves γ homologous to λ .

Recall that for a simple curve λ ,

$$\text{Ext}_\lambda(X) = \sup_\rho \inf_{\gamma \sim \lambda} \frac{\ell_\rho(\gamma)^2}{\text{Area}(\rho)}, \quad (4)$$

where the supremum is over the metrics ρ which are in the conformal class of X , and the inf is over all the curves γ homotopic to λ . Recall that the extremal metric ρ is always the flat metric obtained from a quadratic differential q , and in this metric the entire surface consists of a flat cylinder in which λ is the core curve.

Note that if this quadratic differential q is in fact the square of an Abelian differential then it follows from Theorem 4.2 that

$$\text{Ext}_\lambda(X) = \|c_\lambda\|^2. \quad (5)$$

This is indeed the case for the curves α_i and β_i of Theorem 4.1. This can be used to give an alternative proof of Theorem 4.1.

4.4 The Hodge Norm and the Geodesic Flow.

Let Ω_g denote the space of holomorphic 1-forms on surfaces of genus g . We also denote by g_t denote the geodesic flow on Ω_g (where we square a holomorphic 1-form to get a quadratic differential). Fix a point S in Ω_g ; then S is a pair (X, ω) where ω is a holomorphic 1-form on X . Let $\|\cdot\|_0$ denotes the Hodge norm on the surface $X_0 = X$, and let $\|\cdot\|_t$ denote the Hodge norm on the surface $X_t = \pi(g_t S)$.

The following fundamental result is due to Forni [Fo02, §2]:

Theorem 4.3 *There exists a constant C depending only on the genus, such that for any $\theta \in H^1(X, \mathbb{R})$ and any $t \geq 0$,*

$$\|\theta\|_t \leq Ce^t \|\theta\|_0.$$

If in addition $\langle \theta, \omega \rangle = 0$, and for some compact subset $\mathcal{K}$ of $\mathcal{M}_g$, the geodesic segment $[S, g_t S]$ spends at least half the time in $\pi^{-1}(\mathcal{K})$, then we have

$$\|\theta\|_t \leq Ce^{(1-\alpha)t} \|\theta\|_0,$$

where $\alpha > 0$ depends only on $\mathcal{K}$.

5 Measure in Polar Coordinates

Let $\mathcal{P}(1, \dots, 1) \subset \mathcal{Q}^1 \mathcal{T}_g$ denote the subset where the all zeroes of the quadratic differential are distinct. $\mathcal{P}(1, \dots, 1)$ is called the *principal stratum*. Its complement in $\mathcal{Q}^1 \mathcal{T}_g$ is the *multiple zero locus*.

Recall that $\mathcal{Q}^1 \mathcal{T}_g$ is the space of pairs (M, q) where M is a Riemann surface and q is a holomorphic quadratic differential on M . (We occasionally drop M from the notation and denote points of $\mathcal{Q}^1 \mathcal{T}_g$ by q alone). Let $\tilde{M}$ denote the canonical double cover on which q becomes the square of an Abelian differential. For a ring R' , let $H_{odd}^1(\tilde{M}, R')$ (resp. $H_1^{odd}(\tilde{M}, R')$) denote the part of the first cohomology (resp. homology) of $\tilde{M}$ which is odd under the involution exchanging the two sheets of the double cover. Note that the dimension of $H_1^{odd}(\tilde{M}, \mathbb{R})$ is $h = 6g - 6$.

Fix $x \in \mathcal{T}_g$. Then we have a diffeomorphism $\Phi : S(x) \times \mathbb{R}^+ \rightarrow \mathcal{T}_g$ where $\Phi(q, t) = g_t q$ is the point in $\mathcal{T}$ which is the endpoint of the length t geodesic segment starting at x and tangent to $q \in S(x)$. We can then write

$$d\mathbf{m}(\pi(g_t q)) = \Delta(q, t) ds_x(q) dt. \quad (6)$$

Proposition 5.1 *Let $K_1 \supset K$ be compact subsets of $\mathcal{P}(1, \dots, 1)$, $q \in \mathcal{Q}^1 \mathcal{T}_g$, and suppose q and $g_t q$ both belong to ΓK . Then*

$$|\Delta(q, t)| \leq Ce^{ht}, \quad (7)$$

where C depends only on K . If in addition

$$|\{s \in [0, t] : g_s q \in \Gamma K_1\}| \geq (1/2)t \quad (8)$$

then there exists $\alpha > 0$ so that

$$\Delta(q, t) = e^{ht} \lambda^-(q) \lambda^+(g_t q) + O(e^{(h-\alpha)t}) \quad (9)$$

where α and the implied constant on (9) depend only on K_1 .

Proof. Without loss of generality, $q \in K$. Let $x = \pi(q)$, $y = \pi(g_t q)$. Write $y = \gamma z$ where $\gamma \in \Gamma$ and $z \in K$. Let A be the matrix of γ acting on the odd part of the homology of the orienting double cover $\tilde{M}$ of q (i.e. A is the Kontsevitch-Zorich cocycle.) Let $\langle \cdot, \cdot \rangle$ denote the Hodge inner product, and A^* denote the adjoint of A with respect to $\langle \cdot, \cdot \rangle$. Let $n = h = \dim H_{odd}^1(\tilde{M}, \mathbb{R})$, and let $e_1, \dots, e_n$ be an orthonormal basis for $H_{odd}^1(\tilde{M}, \mathbb{R})$ with respect to $\langle \cdot, \cdot \rangle$, so that

$$A^* A e_i = \lambda_i^2 e_i,$$

where $e^t = \lambda_1 \geq \dots \geq \lambda_n = e^{-t} > 0$. Then $\|A e_i\| = \lambda_i$. Note that e_1 is the imaginary part of the holomorphic 1-form $\tilde{\omega}$ on $\tilde{M}$ for which $\tilde{\omega}^2 = q$, and e_n is the real part of $\tilde{\omega}$. We can read off that $\lambda_1 = e^t$, $\lambda_n = e^{-t}$.

Note since we are assuming $q \in K$, the pair $(\tilde{M}, \tilde{\omega})$ associated to q belongs to a compact subset (depending only on K) of the space of Abelian differentials. Then, by Theorem 4.3,

$$\lambda_2 = O(e^t) \text{ and } \lambda_{n-1}^{-1} = O(e^t), \quad (10)$$

and if (8) holds,

$$\lambda_2 = O(e^{(1-\alpha)t}) \text{ and } \lambda_{n-1}^{-1} = O(e^{(1-\alpha)t}). \quad (11)$$

Since A has determinant 1, the product of the λ_i is 1.

We can identify the tangent space to $\mathcal{Q}^1 \mathcal{T}_g$ with $H_{odd}^1(\tilde{M}, \mathbb{R}^2) = H_{odd}^1(\tilde{M}, \mathbb{R}) \otimes \mathbb{R}^2$ (see e.g. [Fo02, §1] for the details).

Let $e_i^+ = e_i \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $e_i^- = e_i \otimes \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ so that the e_i^+ and e_i^- together form a basis for the tangent space at x . Then $(dg_t)_*(e_i^+) = e^t A e_i^+$, and $(dg_t)_*(e_i^-) = e^{-t} A e_i^-$. (In the above we adopt the convention that A acts on the tensor product $H_{odd}^1(\tilde{M}, \mathbb{R}) \otimes \mathbb{R}^2$ by acting on the $H_{odd}^1(\tilde{M}, \mathbb{R})$ factor, and trivially on the second factor.

Let $u_1 = e_1^+ - e_n^-$. We can complete the set $\{u_1\}$ to a basis $\{u_1, \dots, u_{n-1}\}$ for the tangent space of $S(x)$ at q , so that for $2 \leq i \leq n-1$, $u_i = e_i^+ + w_i$, where w_i is in the span of $\{e_2^-, \dots, e_{n-1}^-\}$. For consistency, we let $w_1 = e_n^-$. Note that the $\|w_i\|$ are bounded since q belongs to a compact set K , and the leaves of $\mathcal{F}^u$ are transverse to the spheres $S(x)$ on K .

We extend the Hodge inner product $\langle \cdot, \cdot \rangle$ to $H_{odd}^1(\tilde{M}, \mathbb{R}^2)$ by declaring that $\langle e_i^+, e_j^- \rangle = 0$ for all i, j . This also extends the Hodge norm. We now estimate

$$\begin{aligned} \|(dg_t)_*(u_1 \wedge \dots \wedge u_n)\|^2 &= \langle u_1 \wedge \dots \wedge u_n, [(dg_t)_*]^* [(dg_t)_*] u_1 \wedge \dots \wedge u_n \rangle \\ &= \sum_P \langle u_1 \wedge \dots \wedge u_n, [(dg_t)_*]^* [(dg_t)_*] \left(\bigwedge_{i \notin P} e_i^+ \wedge \bigwedge_{i \in P} w_i \right) \rangle \\ &= \sum_P \langle u_1 \wedge \dots \wedge u_n, \left(\bigwedge_{i \notin P} e^{2t} \lambda_i^2 e_i^+ \wedge \bigwedge_{i \in P} e^{-2t} A^* A w_i \right) \rangle \end{aligned} \quad (12)$$

where $[(dg_t)_*]^*$ is the adjoint of the linear transformation $(dg_t)_*$, and the sum is over all subsets P of $\{1, \dots, n\}$.

Note that by (10), for $1 \leq i \leq n-1$

$$\|e^{-2t} A^* A w_i\| \leq C \leq C \|e^{2t} \lambda_i^2\|.$$

Thus all the terms in (12) are bounded by $C e^{2t(n-1)} \lambda_1^2 \prod_{i=2}^{n-1} \lambda_i^2 = C e^{nt} = C e^{ht}$. Thus,

$$\|(dg_t)_*(u_1 \wedge \dots \wedge u_n)\| = O(e^{ht}). \quad (13)$$

By definition, $\Delta(q, t)$ is the determinant of the linear transformation $D = (d\pi_y)_*(dg_t)_*$. Since y is in a compact set, the norm of $(d\pi_y)_*$ is bounded. This together with (13) implies (7). If we assume (11), we get for any $1 \leq i \leq n-1$,

$$\|e^{-2t} A^* A w_i\| \leq C e^{-2\alpha t} \leq C e^{-2\alpha} e^{2t} \lambda_i^2,$$

which implies that the contribution of all the terms in (12) to $\Delta(q, t)$ except for the term with $P = \emptyset$ is $O(e^{(h-\alpha)t})$. It remains to evaluate the term with $P = \emptyset$. Dropping the rest of the terms is equivalent to replacing the map $\Phi(q, t)$ by the composition of three maps:

1. Projecting from the sphere $S(x)$ to the leaf of the expanding foliation $\mathcal{F}^u$ passing through q .
2. Flowing by time t .
3. Applying the map π_y to project from the leaf of $\mathcal{F}^u$ through $g_t q$ to Teichmüller space $\mathcal{T}_g$.

The Jacobian of the resulting map is the product of the Jacobians from steps 1), 2) and 3). Step 1) gives a factor of $\lambda^-(q)$, step 2) gives a factor of e^{ht} , and step 3) gives a factor of $\lambda^+(g_t q)$, so one gets formula (9). $\square$

6 Mixing.

Let $\mathcal{U}$ be an open subset of the boundary at infinity of $\mathcal{T}_g$ (which can be identified with the space $\mathcal{PMF}$ of projective measured foliations). For each $x \in \mathcal{T}_g$ we may identify $\mathcal{U}$ with a subset $\mathcal{U}(x)$ of $S(x)$. Let $Sect_{\mathcal{U}}(x) \subset \mathcal{T}_g$ denote the set $\bigcup_{t \geq 0} \pi(g_t \mathcal{U}(x))$, i.e. the sector based at x in the direction $\mathcal{U}$. For a subset $A \subset \mathcal{T}_g$, $Nbhd_r(A)$ denotes the set of points within Teichmüller distance r of A .

Theorem 6.1 *For any $x \in \mathcal{T}_g$ and any $\epsilon > 0$ there exists an open subset $\mathcal{U}$ of $S(x)$ containing the intersection of $S(x)$ with the multiple zero locus such that for any compact $\mathcal{K} \subset \mathcal{T}_g$*

$$\limsup_{R \rightarrow \infty} e^{-hR} \mathbf{m}(Nbhd_1(B_R(x) \cap Sect_{\mathcal{U}}(x) \cap \Gamma \mathcal{K})) \leq \epsilon \quad (14)$$

Let $\mathcal{K}$ be a compact subset of $\mathcal{T}_g$, and let K' be a compact subset of the principal stratum $\mathcal{P}(1, \dots, 1) \subset \mathcal{Q}^1 \mathcal{T}_g$. Let $B_R(x, \mathcal{K}, K')$ denote the set of points $y \in \Gamma \mathcal{K}$ such that $d(x, y) < R$, and the geodesic from x to y spends at least half the time outside $\Gamma K'$.

The following theorem will also be proved in §9:

Theorem 6.2 *For $\epsilon > 0$ and any compact $\mathcal{K} \subset \mathcal{T}_g$ there exists compact $K' \subset \mathcal{P}(1, \dots, 1) \subset \mathcal{Q}^1 \mathcal{T}_g$ such that for any $x \in \mathcal{K}$,*

$$\limsup_{R \rightarrow \infty} e^{-hR} \mathbf{m}(Nbhd_1(B_R(x, \mathcal{K}, K')))) \leq \epsilon \quad (15)$$

Let ψ be a nonnegative compactly supported continuous function on $\mathcal{M}_g$, which we extend to a function $\psi : \mathcal{Q}^1 \mathcal{M}_g \rightarrow \mathbb{R}$ by making it constant on each sphere $S(x)$. We can consider ψ to be a Γ -periodic function on $\mathcal{T}_g$ (or $\mathcal{Q}^1 \mathcal{T}_g$). Let ϕ be another such function.

Proposition 6.3 (Mean Ergodic Theorem) *We have*

$$\lim_{R \rightarrow \infty} h e^{-hR} \int_{\mathcal{M}_g} \phi(x) \left(\int_{B_R(x)} \psi(y) d\mathbf{m}(y) \right) d\mathbf{m}(x) = \int_{\mathcal{M}_g} \phi(x) \Lambda(x) d\mathbf{m}(x) \int_{\mathcal{M}_g} \psi(y) \Lambda(y) d\mathbf{m}(y). \quad (16)$$

Proof. Suppose $\epsilon > 0$ is given. Let $\mathcal{K}$ be the union of the supports of ϕ and ψ . Without loss of generality, we may assume that the support of ϕ is small enough so that there exists an open $\mathcal{U} \subset \partial\mathcal{T}_g$ such that for each x in the support of ϕ , $\mathcal{U}(x)$ contains the intersection of $S(x)$ with the multiple zero locus, and also (14) holds.

We break up the integral over $B_R(x)$ in (16) into two parts: the first over $B_R(x) \cap \text{Sect}_{\mathcal{U}}(x) \cap \Gamma\mathcal{K}$ and the second over the complement. The first is bounded by $C_1\epsilon$ in view of Theorem 6.2.

Let K' be as in Theorem 6.2. We use polar coordinates for the integral over $B_R(x)$. We get,

$$\int_{\mathcal{M}_g} \phi(x) \left(\int_{B_R(x)} \psi(y) d\mathbf{m}(y) \right) d\mathbf{m}(x) = \int_{\mathcal{M}_g} \phi(x) \left(\int_0^R \int_{S(x)} \Delta(q, t) \psi(g_t q) ds_x(q) dt \right) d\mathbf{m}(x). \quad (17)$$

In view of (15), we may assume, (up to error of size $C\epsilon e^{hR}$) that $q \in S(x)$ belongs to the compact set $S(x) \setminus \mathcal{U}(x)$, which is away from the multiple zero locus. Also note that the left-hand-side of (17) is symmetric in x and y (up to interchanging the functions ϕ and ψ). So we may also assume that $g_t q$ belongs to a fixed compact set which is bounded away from the multiple zero locus. Also in view of (15), we may assume that (8) holds. (Again the contribution of the part of the integral where this assumption is violated is bounded by $C\epsilon e^{hR}$.)

For the part of the integral where none of the three assumptions are violated, we are finally in a position to use Proposition 5.1 to replace $\Delta(q, t)$ by $e^{ht} \lambda^-(q) \lambda^+(g_t q)$. Then, the right hand side of (17) becomes

$$\begin{aligned} \int_0^R e^{ht} \int_{\mathcal{T}_g} \phi(x) \int_{S(x)} \lambda^-(q) \psi(g_t q) \lambda^+(g_t q) ds_x(q) d\mathbf{m}(x) dt = \\ \int_0^R e^{ht} \int_{\mathcal{Q}^1 \mathcal{M}_g} \phi(q) \lambda^-(q) \psi(g_t q) \lambda^+(g_t q) d\mu(q) dt, \end{aligned} \quad (18)$$

where we used the fact that ϕ is constant on each sphere $S(x)$ and the formula

$$d\mu(q) = d\mathbf{m}(\pi(q)) d_{s_{\pi(q)}}(q).$$

Now we can apply the mixing property of the geodesic flow to the functions $\phi \lambda^-$ and $\psi \lambda^+$. The right-hand side of (18) is then asymptotic to

$$\frac{e^{hR}}{h} \int_{\mathcal{Q}^1 \mathcal{M}_g} \phi(q) \lambda^-(q) d\mu(q) \int_{\mathcal{Q}^1 \mathcal{M}_g} \psi(q) \lambda^+(q) d\mu(q) = \frac{e^{hR}}{h} \int_{\mathcal{M}_g} \phi(x) \Lambda(x) d\mathbf{m}(x) \int_{\mathcal{M}_g} \psi(y) \Lambda(y) d\mathbf{m}(y)$$

So the right hand side of (16) is equal to the left hand side of (16) up to an error which is bounded by $C\epsilon$, where C depends only on ϕ and ψ . Since $\epsilon > 0$ is arbitrary, the theorem follows. $\square$

The proof of Proposition 6.3 also shows that for any open $\mathcal{U} \subset S(x)$,

$$\begin{aligned} \lim_{R \rightarrow \infty} h e^{-hR} \int_{\mathcal{M}_g} \phi(x) \left(\int_{B_R(x) \cap \text{Sect}_{\mathcal{U}}(x)} \psi(y) d\mathbf{m}(y) \right) d\mathbf{m}(x) = \\ \int_{\mathcal{M}_g} \phi(x) \left(\int_{\mathcal{U}} \lambda^-(q) ds_x(q) \right) d\mathbf{m}(x) \int_{\mathcal{M}_g} \psi(y) \Lambda(y) d\mathbf{m}(y). \end{aligned} \quad (19)$$

7 Counting

Proof of Theorem 1.1. Let $F_R(x, y)$ denote the cardinality of the intersection of $B_R(x)$ with Γy . Let ϕ and ψ be a negative compactly supported continuous functions on $\mathcal{M}_g$. Note that

$$\int_{\mathcal{M}_g} F_R(x, y) \psi(y) d\mathbf{m}(y) = \int_{B_R(x)} \psi(y) d\mathbf{m}(y).$$

Hence, by Proposition 6.3, as $R \rightarrow \infty$,

$$e^{-hR} \int_{\mathcal{M}_g} \phi(x) \int_{\mathcal{M}_g} F_R(x, y) \psi(y) d\mathbf{m}(y) d\mathbf{m}(x) \rightarrow \frac{1}{h} \int_{\mathcal{M}_g} \phi(x) \Lambda(x) d\mathbf{m}(x) \int_{\mathcal{M}_g} \psi(y) \Lambda(y) d\mathbf{m}(y),$$

i.e. $e^{-hR} F_R(x, y)$ converges weakly to $\frac{1}{h} \Lambda(x) \Lambda(y)$. We would like to show that the convergence is pointwise on compact sets.

Let $\epsilon > 0$ be arbitrary. Let ψ be supported on a ball of radius ϵ around y (in the Teichmüller distance), and satisfy $\int_{\mathcal{M}_g} \psi d\mathbf{m} = 1$. Let ϕ be supported on a ball of radius ϵ around x , with $\int_{\mathcal{M}_g} \phi d\mathbf{m} = 1$. By the triangle inequality,

$$\int_{\mathcal{M}_g} \phi(x) \int_{B_{R-3\epsilon}(x)} \psi(y) d\mathbf{m}(y) d\mathbf{m}(x) \leq F_R(x, y) \leq \int_{\mathcal{M}_g} \phi(x) \int_{B_{R+3\epsilon}(x)} \psi(y) d\mathbf{m}(y) d\mathbf{m}(x).$$

After multiplying both sides by $h e^{-hR}$ and taking $R \rightarrow \infty$, we get, after applying Proposition 6.3,

$$\begin{aligned} e^{-3h\epsilon} \int_{\mathcal{M}_g} \phi(x) \Lambda(x) d\mathbf{m}(x) \int_{\mathcal{M}_g} \psi(y) \Lambda(y) d\mathbf{m}(y) &\leq \liminf_{R \rightarrow \infty} h e^{-hR} F_R(x, y) \leq \\ &\leq \limsup_{R \rightarrow \infty} h e^{-hR} F_R(x, y) \leq e^{3h\epsilon} \int_{\mathcal{M}_g} \phi(x) \Lambda(x) d\mathbf{m}(x) \int_{\mathcal{M}_g} \psi(y) \Lambda(y) d\mathbf{m}(y). \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, and Λ is uniformly continuous on compact sets, we get

$$\lim_{R \rightarrow \infty} h e^{-hR} F_R(x, y) = \Lambda(x) \Lambda(y)$$

as required. □

In fact, the same argument using (19) instead of Proposition 6.3 yields the following:

Theorem 7.1 Suppose $x \in \mathcal{T}_g$, and $\mathcal{U}$ is an open subset of the boundary at infinity of $\mathcal{T}_g$. Then for any $y \in \mathcal{T}_g$, we have, as $R \rightarrow \infty$,

$$|B_R(x) \cap \text{Sect}_{\mathcal{U}}(x) \cap \Gamma y| \sim \frac{1}{h} e^{hR} \Lambda(y) \int_{\mathcal{U}} \lambda^-(q) ds_x(q).$$

We also obtain the following:

Theorem 7.2 Suppose $x \in \mathcal{T}_g$, and $\mathcal{U}$ and $\mathcal{V}$ are open subsets of the boundary at infinity of $\mathcal{T}_g$. Then for any $y \in \mathcal{T}_g$, we have, as $R \rightarrow \infty$,

$$|\{\gamma \in \Gamma : \gamma y \in B_R(x) \cap \text{Sect}_{\mathcal{U}}(x) \text{ and } \gamma^{-1}x \in \text{Sect}_{\mathcal{V}}(y)\}| \sim \frac{1}{h} e^{hR} \int_{\mathcal{U}} \lambda^-(q) ds_x(q) \int_{\mathcal{V}} \lambda^+(q) ds_y(q).$$

The proof of Theorem 7.2 is very similar to that of Theorem 7.1 except that we do not assume that the function $\psi : \mathcal{Q}^1 \mathcal{M}_g \rightarrow \mathbb{R}$ is the pullback under π of a function from $\mathcal{M}_g \rightarrow \mathbb{R}$. The details are left to the reader.

Patterson-Sullivan Measure. Let the measure ν_x on $\mathcal{PMF}$ be the image of $\lambda^-(q) ds_x(q)$ under the identification of $S(x)$ and $\mathcal{PMF}$. Theorem 7.1 implies that ν_x is the *Patterson-Sullivan* measure. This measure coincides with the Thurston measure on $\mathcal{PMF}$

$$\nu_x(U) = \mu_{Th}(\{\eta \mid [\eta] \in U, \sqrt{\text{Ext}_{\eta}(X)} \leq 1\}).$$

By definition, for $x, y \in \mathcal{T}_g$, $\xi \in \mathcal{MF}$, we have

$$\frac{d\nu_x}{d\nu_y}([\xi]) = \left(\frac{\sqrt{\text{Ext}_{\xi}(y)}}{\sqrt{\text{Ext}_{\xi}(x)}} \right)^{6g-6},$$

where $[\xi]$ denotes the image of ξ in $\mathcal{PMF}$; note that the right-hand side only depends on $[\xi]$.

- Given $\xi \in \mathcal{PMF}$, let $\beta_{\xi} : \mathcal{T}_g \times \mathcal{T}_g \rightarrow \mathbb{R}$ be the cocycle defined by

$$\beta_{\xi}(x, y) = \log \left(\frac{\sqrt{\text{Ext}_{\xi}(x)}}{\sqrt{\text{Ext}_{\xi}(y)}} \right).$$

In our setting β_{ξ} plays the role of the Busemann cocycle for the Teichmüller metric, and (formally) the family $\{\nu_x\}_{x \in \mathcal{T}_g}$ of finite measures on $\mathcal{PMF}$ is a family of *conformal densities* of dimension $\delta = 6g - 6$ for the cocycle β ; i.e, for any $x, y \in \mathcal{T}_g$ and almost every $\xi \in \mathcal{PMF}$ we have

$$\frac{d\nu_x}{d\nu_y}([\xi]) = \exp(\delta \beta_{\xi}(y, x)), \text{ and } \nu_{g \cdot x}(g \cdot U) = \nu_x(U),$$

where $U \subset \mathcal{PMF}$, and g is an element of the mapping class group.

Moreover, by the measure classification result of [LM], $\{\nu_x\}_{x \in \mathcal{T}_g}$ is the unique conformal density for the action of the mapping class group on $\mathcal{T}_g$ up to scale; the only possibility for δ is $6g - 6$.

- Given $x \in \mathcal{T}_g$, and transverse minimal foliations $\xi, \eta \in \mathcal{MF}$, define

$$\beta(x, [\xi], [\eta]) = \left(\frac{\sqrt{\text{Ext}_\xi(x) \text{Ext}_\eta(x)}}{A(\xi, \eta)} \right)^{6g-6},$$

where $A(\xi, \eta)$ is the area of the quadratic differential with horizontal foliation ξ , and vertical foliation η . Note that the right-hand side only depends on $[\xi], [\eta]$.

Now introduce a measure $\mathbb{P}$ on the product $\mathcal{PMF} \times \mathcal{PMF}$ by the formula

$$\mathbb{P} = \beta(x, [\xi], [\eta]) \nu_x([\xi]) \nu_x([\eta]). \quad (20)$$

Observe that the right-hand side does not depend on x and is invariant under the diagonal action of the mapping class group.

The space $\mathcal{Q}^1 \mathcal{T}_g$ of quadratic differentials fibres over $\mathcal{PMF} \times \mathcal{PMF}$ (with fibre $\mathbb{R}$) and the measure $\mathbb{P}$ on $\mathcal{PMF} \times \mathcal{PMF}$ lifts to (a multiple of) the Lebesgue measure on $\mathcal{Q}^1 \mathcal{T}_g$ (by construction, it is invariant and belongs to the Lebesgue measure class — but this determines the measure uniquely).

These constructions are similar to the ones due to Kaimanovich in [Ka90a] and [Ka90b].

8 The Hodge Norm and Quadratic Differentials

8.1 The Hodge Norm and the Flat Metric.

Theorem 4.1 allows us to estimate the Hodge norm in terms of the hyperbolic metric. In this subsection we state a lemma which allows us to estimate the Hodge norm in terms of the flat metric. This is done by first estimating the hyperbolic metric in terms of the flat metric, and then using Theorem 4.1.

Let q be a holomorphic quadratic differential, and let ℓ_q denote length in the flat metric defined by q . For each $\epsilon > 0$ let $K(\epsilon)$ denote the complex generated by all saddle connections shorter than ϵ (see [EM01, §6]). Recall that $K(\epsilon)$ contains all saddle connections shorter than ϵ , and that $\ell_q(\partial K(\epsilon)) = O(\epsilon)$, where the implied constant depends only on the genus. (Also if C is a cylinder whose boundary is shorter than ϵ , we include C in $K(\epsilon)$). Let $\epsilon_1 < \dots < \epsilon_n$ denote the values of ϵ where $K(\epsilon)$ changes. (Note that n is bounded in terms of the genus). Pick a constant $C \gg 1$ and drop all ϵ_i such that $\epsilon_i > \epsilon_{i+1}/C$. After renumbering, we obtain new ϵ_i $1 \leq i \leq m$, with $\epsilon_i < \epsilon_{i+1}/C$. To simplify notation, we let $\epsilon_{m+1} = 1$. For $1 \leq i \leq m$, let K_i denote the complex $K(\epsilon_i)$.

Lemma 8.1 (Rafi, [Ra05]) *Let q be a holomorphic quadratic differential on a surface S , and let σ be the hyperbolic metric on S in the same conformal class as q . Choose $C \gg 1$, and let the ϵ_i , $1 \leq i \leq m$ and the complexes K_i be as in the previous paragraph. Then*

- There exists a constant ϵ_0 depending only on C and the genus (with $\epsilon_0 \rightarrow 0$ as $C \rightarrow \infty$) such that any simple closed curve α on S with the hyperbolic length $\ell_\sigma(\alpha) < \epsilon_0$ is homotopic to a connected component of the boundary of one of the K_i .*
- Let γ be a connected component of ∂K_i . If γ is not a core curve of a flat annulus, then $\ell_\sigma(\gamma) \geq C |\log \epsilon_i|$, where C is a constant depending only on the genus.*

8.2 The Modified Hodge Norm and Quadratic Differentials

Theorem 4.3 gives a partial hyperbolicity property of the geodesic flow on spaces of Abelian differentials. In our applications, we need a similar property for the spaces $\mathcal{Q}^1\mathcal{T}_g$ of quadratic differentials. A standard construction is, given $M \in \mathcal{T}_g$ and q a holomorphic quadratic differential on M , is to pass to the double cover on which the foliation defined by q is orientable. This yields a surface $\tilde{M}$ with an holomorphic Abelian differential ω . However, a major difficulty is the following: even if q belongs to a compact subset of $\mathcal{Q}^1\mathcal{T}_g$, the complex structure on $\tilde{M}$ may have very short closed curves in the hyperbolic metric. This may occur since even if one restricts q to compact subsets of Teichmüller space, the flat structure defined by q may have arbitrarily short saddle connections (connecting distinct zeroes). Such a saddle connection may lift to a very short loop when one takes a double cover. Other more complicated types of degeneration are also possible. However, the following simple observation is key to our approach:

Lemma 8.2 *For every compact subset $\mathcal{K} \subset \mathcal{M}_g$ there exists a constant $\epsilon_0 > 0$ depending only on $\mathcal{K}$ such that for any $q \in \pi^{-1}(\Gamma\mathcal{K})$, the flat structure associated to the holomorphic quadratic differential on the orienting cover of q has no closed trajectories of Euclidean length less than ϵ which are part of a flat cylinder.*

Together with Lemma 8.1 (b), Lemma 8.2 will allow us to estimate the hyperbolic length of short curves on $\tilde{M}$.

Proof of Lemma 8.2. Given the set $\mathcal{K}$, there exists a constant $\epsilon_0 > 0$ such that any loop of length less than ϵ_0 in the flat metric is contractible. Now suppose $q \in \pi^{-1}(\mathcal{K})$, and γ is a trajectory on the associated orienting double cover which has length less than ϵ_0 , and is part of a flat cylinder C . Let γ_1 be the projection of γ to the original flat structure defined by q . Then γ_1 must also be a part of a flat cylinder, and the length of γ_1 must be at most ϵ_0 . Then, from the definition of ϵ_0 it follows that γ_1 must be contractible. But this is impossible since the curvature of γ_1 is zero, and we are not allowing q to have poles. $\square$

Short bases. Suppose $(X, \omega) \in \Omega_g$. Fix $\epsilon_1 < \epsilon_0$ and let $\alpha_1, \dots, \alpha_k$ be the curves with hyperbolic length less than ϵ_1 on X . For $1 \leq i \leq k$, let β_i be the shortest curve in the flat metric defined by ω with $i(\alpha_i, \beta_i) = 1$. Let γ_r , $1 \leq r \leq 2g - 2k$ be moderate length curves on X so that the α_j , β_j and γ_j are a symplectic basis $\mathcal{S}$ for $H_1(X, \mathbb{R})$. We will call such a basis *short*.

The functions $\chi_i(\omega)$ and the modified Hodge norm. We would like to use the Hodge norm on double covers of surfaces in $\mathcal{Q}^1\mathcal{T}_g$. One difficulty is that if one takes the double cover of some quadratic differential in the multiple zero locus, the Hodge norm in some directions tangent to the multiple zero locus might vanish. As a consequence, if we use the Hodge norm to define a “Hodge distance” on $\mathcal{Q}^1\mathcal{T}_g$, then we find that some balls are not compact, and the resulting “distance” does not separate points on the multiple zero locus.

We now define a modification of the Hodge norm in order to avoid these problems. The modified norm is a norm on the tangent space to the space of pairs (X, ω) where X is a Riemann surface and ω is a holomorphic 1-form on X . Unlike the Hodge norm, the modified Hodge norm will depend not only on the complex structure on X but also on the choice of a holomorphic 1-form ω on X . Let $\{\alpha_j, \beta_j, \gamma_r\}_{1 \leq j \leq k, 1 \leq r \leq 2g-2k}$ be a short basis for (X, ω) . For $1 \leq i \leq k$ we define $\chi_i(\omega)$ to be 0 if α_i has a flat annulus in the flat metric defined by ω and 1 otherwise.

We can write any $\theta \in H^1(X, \mathbb{R})$ as

$$\theta = \sum_{i=1}^k a_i (*c_{\alpha_i}) + \sum_{i=1}^k b_i \ell_\sigma(\alpha_i)^{1/2} (*c_{\beta_i}) + \sum_{r=1}^{2g-2k} u_r (*c_{\gamma_r}). \quad (21)$$

(Recall that σ denotes the hyperbolic metric in the conformal class of X , and for a curve α on X , $\ell_\sigma(\alpha)$ denotes the length of α in the metric σ). We then define

$$\|\theta\|' = \|\theta\| + \left(\sum_{i=1}^k \chi_i(\omega) |a_i| + \sum_{i=1}^k |b_i| + \sum_{r=1}^{2g-2k} |u_r| \right) \quad (22)$$

From (22) we have for $1 \leq i \leq k$,

$$\|*c_{\alpha_i}\|' \approx 1, \quad (23)$$

as long as α_i has no flat annulus in the metric defined by ω . Similarly, from (2) we have

$$\|*c_{\beta_i}\|' \approx \|*c_{\beta_i}\| \approx \frac{1}{\ell_\sigma(\alpha_i)^{1/2}} \quad (24)$$

In addition, in view of Theorem 4.1, if γ is any other moderate length curve on X , $\|*c_\gamma\|' \approx \|*c_\gamma\| = O(1)$. Thus, if $\mathcal{B}$ is a short basis associated to ω , then for any $\gamma \in \mathcal{B}$,

$$\text{Ext}_\gamma(\omega)^{1/2} \approx \|*c_\gamma\| \leq \|*c_\gamma\|' \quad (25)$$

(cf. (5)). (By $\text{Ext}_\gamma(\omega)$ we mean the extremal length of γ in the conformal structure defined by ω).

Remark. From the construction, we see that the modified Hodge norm is greater than the Hodge norm. Also, if the flat length of shortest curve in the flat metric defined by ω is greater than ϵ_1 , then for any cohomology class λ , for some C depending on ϵ_1 and the genus,

$$\|\lambda\|' \leq C\|\lambda\|, \quad (26)$$

i.e. the modified Hodge norm is within a multiplicative constant of the Hodge norm.

8.3 Non-expansion of the modified Hodge norm.

In this subsection we will prove a weak version of Theorem 4.3 for the modified Hodge norm.

Let $\|\cdot\|'_0$ denote the modified Hodge norm on the surface $S_0 = S = (X_0, \omega_0)$, and let $\|\cdot\|'_t$ denote the modified Hodge norm on the surface $S_t = g_t S = (X_t, \omega_t)$.

Theorem 8.3 *There exists a constant C depending only on the genus and on ϵ_1 , such that for any $\theta \in H^1(X, \mathbb{R})$ and any $t \geq 0$,*

$$\|\theta\|'_t \leq C e^t \|\theta\|'_0. \quad (27)$$

The proof of Theorem 8.3 is based on the following lemma:

Lemma 8.4 (Rafi) *Suppose $(X_0, \omega_0) \in \Omega_g$, and write (X_t, ω_t) for $g_t(X_0, \omega_0)$. Suppose $\mathcal{B}$ is a short basis in the flat structure defined by ω_0 , and $\mathcal{B}'$ is a short basis in the flat structure defined by ω_t . Suppose α is a curve of hyperbolic length $< \epsilon_1$ on both X_0 and X_t (so that in particular $\alpha \in \mathcal{B}$ and*

$\alpha \in \mathcal{B}'$), and that α has no flat annulus. Let $\beta' \in \mathcal{B}'$ denote the curve with $i(\alpha, \beta') = 1$. Then, for any $\gamma \in \mathcal{B}$, we have

$$i(\gamma, \beta') \leq C e^t \text{Ext}_\gamma(X_0)^{1/2},$$

where $i(\cdot, \cdot)$ denotes the intersection number.

Proof. First we claim that $i(\gamma, \beta') \leq i(\gamma, P)$, where P is the shortest pants decomposition of X_t . This is because, either γ is disjoint from α or if it intersects α , the relative twisting of γ and β' is bounded (see [Ra07, §4]). But this shortest pants decomposition has extremal length less than e^{2t} in X_0 . Using [Mi93, Lemma 5.1] we have

$$i(\gamma, P)^2 \leq C^2 \text{Ext}_\gamma(X_0) \text{Ext}_P(X_0) \leq C^2 e^{2t} \text{Ext}_\gamma(X_0).$$

This finishes the proof. $\square$

Proof of Theorem 8.3. Let $\alpha'_1, \dots, \alpha'_k$ be the curves with hyperbolic length less than ϵ_1 on $X_t = \pi(S_t)$. Let $\beta'_1, \dots, \beta'_k$ be the shortest curves in the Euclidean metric on S_t such that $i(\alpha_r, \beta_j) = \delta_{rj}$. Let γ'_r , $1 \leq r \leq 2g - k$ be moderate length curves on X_t so that the α_j , β'_j and γ'_j are a symplectic basis $\mathcal{S}'$ for $H_1(X, \mathbb{R})$. Thus $\mathcal{S}'$ is a short basis. For any $\theta \in H^1(X, \mathbb{R})$, we may write

$$\theta = \sum_{i=1}^k a_i (*c_{\alpha'_i}) + \sum_{i=1}^k b_i \ell_\sigma(\alpha'_i)^{1/2} (*c_{\beta'_i}) + \sum_{r=1}^{2g-k} u_i (*c_{\gamma'_i}). \quad (28)$$

In view of (22) and Theorem 4.3, it is enough to show that

$$\sum_{i=1}^k \chi_i(\omega_t) |a_i| + \sum_{i=1}^k |b_i| + \sum_{r=1}^{2g-k} |u_i| \leq C e^t \|\theta\|'_0 \quad (29)$$

We have

$$b_i = \frac{1}{\ell_\sigma(\alpha'_i)^{1/2}} \langle c_{\alpha'_i}, \theta \rangle_t \leq \frac{1}{\ell_\sigma(\alpha'_i)^{1/2}} \|c_{\alpha'_i}\|_t \|\theta\|_t \leq \|\theta\|_t \leq C e^t \|\theta\|_0 \leq C e^t \|\theta\|'_0$$

Also let $\delta'_i \in \mathcal{S}'$ be the element with $i(\gamma'_i, \delta'_i) = 1$. Then,

$$|u_i| = \langle c_{\delta'_i}, \theta \rangle_t \leq \|c_{\delta'_i}\|_t \|\theta\|_t \leq C_1 \|\theta\|_t \leq C_2 e^t \|\theta\|_0 \leq C_3 e^t \|\theta\|'_0$$

It remains to bound $|a_i|$. We may assume that α_i has length less than ϵ_1 on X_0 as well. (If α_i is longer then ϵ_1 on X_0 , we can put in an intermediate point X'_0 where α_i has length $\approx \epsilon_1$.)

Let $\mathcal{S} = \{\alpha_i, \beta_i, \gamma_i\}$ be a short basis for S_0 . It is enough to prove (29) assuming $\theta = *c_\lambda$ for some $\lambda \in \mathcal{S}$. Then,

$$a_i = \langle *c_{\beta'_i}, c_\lambda \rangle = I(\lambda, \beta'_i)$$

Since the algebraic intersection number $I(\cdot, \cdot)$ is bounded by the geometric intersection number $i(\cdot, \cdot)$, we have

$$|a_i| = |I(\lambda, \beta'_i)| \leq i(\lambda, \beta'_i) \leq C e^t \text{Ext}_\lambda(X_0)^{1/2} \leq C e^t \|*c_\lambda\|',$$

where we have used Lemma 8.4 and (25). $\square$

8.4 The Euclidean, Teichmüller and Hodge Distances

Note that we can locally identify a leaf of $\mathcal{F}^s$ (or $\mathcal{F}^u$) with a subspace of $H^1(\tilde{M}, \mathbb{R})$, where $\tilde{M}$ is the double cover of $M \in \mathcal{T}_g$. If γ is a map from $[0, r]$ into some leaf of $\mathcal{F}^{ss}$, then we define the (modified) Hodge length $\ell(\gamma)$ of γ as $\int_0^r \|\gamma'(t)\|' dt$, where $\|\cdot\|'$ is the modified Hodge norm. If q and q' belong to the same leaf of $\mathcal{F}^{ss}$, then we define the Hodge distance $d_H(q, q')$ to be the infimum of $\ell(\gamma)$ where γ varies over paths connecting q and q' and staying in the leaf of $\mathcal{F}^{ss}$ containing q and q' . We make the same definition if q and q' are on the same leaf of $\mathcal{F}^{uu}$.

In what follows $\mathcal{K}$ is a compact subset of $\mathcal{T}_g$, and all implied constants depend on $\mathcal{K}$. Given the set $\mathcal{K}$, there exists a constant $\epsilon_0 > 0$ such that for any $q \in \pi^{-1}(\mathcal{K})$ any loop of length less than ϵ_0 in the flat metric given by q is contractible.

For each $q \in \pi^{-1}(\mathcal{K})$ there exists a canonical (marked) Delaunay triangulation of the flat metric given by q (see [MasSm91, §4]). Since there are only finitely many combinatorial types of (unmarked) triangulations on a genus g surface, and the mapping class group Γ acts properly discontinuously, there are only finitely many combinatorial types α of marked Delaunay triangulations on $\pi^{-1}(\mathcal{K})$. Call this set I . We have

$$\pi^{-1}(\mathcal{K}) = \bigsqcup_{\alpha \in I} W_\alpha,$$

where W_α is the subset of $\pi^{-1}(\mathcal{K})$ where the combinatorial type of the marked Delaunay triangulation is given by α . Let $J \subset I$ denote the subset such that for $\alpha \in J$, W_α is relatively open in $\pi^{-1}(\mathcal{K})$. Then we may write

$$\pi^{-1}(\mathcal{K}) = \bigsqcup_{\alpha \in J} W'_\alpha,$$

where $W_\alpha \subseteq W'_\alpha \subseteq \overline{W_\alpha}$. If we chose a basis $\mathcal{S}$ for $H_1^{odd}(\tilde{M}, \mathbb{Z})$, then we have a period map $\Phi_{\mathcal{S}} : \mathcal{Q}^1 \mathcal{T}_g \rightarrow \mathbb{C}^h$ given by integrating the canonical square root of q along the chosen basis. Recall that this is a local coordinate system for $\mathcal{Q}^1 \mathcal{T}_g$. In fact the following holds:

Lemma 8.5 *For each $\alpha \in J$ there exists a basis $\mathcal{S}_\alpha$ of $H_1^{odd}(\tilde{M}, \mathbb{Z})$ consisting of lifts of edges of the Delaunay triangulation such that the map $\Phi_\alpha = \Phi_{\mathcal{S}_\alpha}$ is a coordinate system on W'_α .*

We can now define a “Euclidean norm” on the tangent space of $\mathcal{Q}^1 \mathcal{T}_g$ as follows: if v is a tangent vector based at the point $q \in W_\alpha$ we define

$$\|v\|_E = |\Phi_\alpha^*(v)|.$$

where $|\cdot|$ is the standard norm on $\mathbb{C}^h$. The norm $\|\cdot\|_E$ depends on the choices of the $\mathcal{S}_\alpha$ and the W'_α ; however if $\|\cdot\|'_E$ is the Euclidean norm obtained from a different choices, then the ratio of $\|\cdot\|_E$ and $\|\cdot\|'_E$ is bounded by a constant depending only on $\mathcal{K}$.

The Euclidean Distance. We define the Euclidean length of a path $\gamma : [0, r] \rightarrow \mathcal{Q}^1 \mathcal{T}_g$ to be $\int_0^r \|\gamma'(t)\|_E dt$. We define the Euclidean distance $d_E(q, q')$ to be the infimum of the Euclidean length of paths connecting q and q' . Locally, up to a multiplicative constant, $d_E(q, q') = |\Phi_\alpha(q) - \Phi_\alpha(q')|$.

We denote the Teichmüller distance on $\mathcal{T}_g$ by $d_{\mathcal{T}}$. For $p \in \mathcal{Q}^1 \mathcal{T}_g$ and $\epsilon > 0$ let $B_E(p, \epsilon)$ denote the ball in the Euclidean metric of radius ϵ centered at p , and let $B_{\mathcal{T}}(\pi(p), \epsilon)$ denote the analogous ball in the Teichmüller metric. We note the following:

Lemma 8.6 *There exist continuous functions $f_i : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $f_i(r) \rightarrow 0$ as $r \rightarrow 0$ such that for any two points p_1 and p_2 in $\pi^{-1}(\mathcal{K})$ on the same leaf of $\mathcal{F}^{ss}$, we have*

$$f_1(d_E(p_1, p_2)) \leq d_{\mathcal{T}}(\pi(p_1), \pi(p_2)) \leq f_2(d_E(p_1, p_2))$$

Proof. This is just a compactness argument. $\square$

The following theorem is the main result of this subsection:

Theorem 8.7 *There exists constants $0 < c_1 < c_2$, both depending only on $\mathcal{K}$, such that for any $q_1, q_2 \in \pi^{-1}(\mathcal{K})$ on the same leaf of $\mathcal{F}^{uu}$ with $d_E(q_1, q_2) < 1$ we have:*

$$c_1 d_E(q_1, q_2) \leq d_H(q_1, q_2) \leq c_2 d_E(q_1, q_2) |\log d_E(q_1, q_2)|^{1/2}.$$

Thus, in particular, on compact sets, the modified Hodge metric is equivalent to the Euclidean metric, and then, in view of Lemma 8.6, the modified Hodge metric on the restriction of a leaf of $\mathcal{F}^{uu}$ to $\pi^{-1}(\mathcal{K})$ is equivalent to the Teichmüller metric.

Note that the lower bound in Theorem 8.7 is clear since the modified Hodge norm is always bigger than the Euclidean norm, see (23) and (24). The rest of this subsection will consist of the proof of the upper bound in Theorem 8.7. For $q \in \mathcal{Q}^1 \mathcal{T}_g$, let $\ell(q)$ denote the length of the shortest saddle connection in the flat metric defined by q .

We will use the following standard fact about Delaunay triangulations:

Lemma 8.8 *Suppose T is the Delaunay triangulation of a surface in $R \in \mathcal{Q}^1 \mathcal{T}_g$. Let $\ell(R)$ denote the length of the shortest saddle connection in R . Then any saddle connection in R of length at most $\sqrt{2}\ell(R)$ is an edge of T .*

Proof. Let $z_1, \dots, z_n$ denote the zeroes of R (i.e. the conical points in the flat metric). For each z_i , let the *Voronoi polygon* V_i denote the set of points of R that are closer to z_i than to any other z_j , $j \neq i$.

Recall that the Delaunay triangulation T of R is dual to the Voronoi diagram in the sense that z_i and z_j are connected by an edge of T if V_i and V_j share a common edge (and also under some conditions if V_i and V_j share a common vertex).

Let e be a saddle connection in R , connecting z_i and z_j . (See Figure 1). Let m be the midpoint of e . Let k be such that $d(z_k, m)$ is minimal. Suppose e is *not* an edge of the Delaunay triangulation T . Then $k \neq i, j$ and $d(m, z_k) \leq d(m, z_i) = \frac{1}{2}d(z_i, z_j)$. We may assume that the angle at m between the segments $\overline{mz_i}$ and $\overline{mz_k}$ is smaller than $\pi/2$ (otherwise replace z_i by z_j). Let p be the point on the segment $\overline{mz_i}$ such that $d(p, m) = d(p, z_k)$. Now consider the isosceles acute triangle τ whose vertices are m, p and z_k . This triangle cannot contain any zeroes (or else z_k would not be the saddle connection closest to m). Therefore, $d(p, z_k) \leq \sqrt{2}d(m, z_k)$. Hence,

$$\begin{aligned} \ell(R) &\leq d(z_i, z_k) \leq d(z_i, p) + d(p, z_k) \leq (d(z_i, m) - d(m, z_k)) + \sqrt{2}d(m, z_k) \\ &= \frac{1}{2}d(z_i, z_j) + (\sqrt{2} - 1)d(m, z_k) \leq \frac{1}{2}d(z_i, z_j) + \frac{(\sqrt{2} - 1)}{2}d(z_i, z_j) = \frac{\sqrt{2}}{2}d(z_i, z_j). \end{aligned}$$

Thus, if e is not an edge of the Delaunay triangulation, then $\ell(e) \geq \sqrt{2}\ell(R)$. $\square$

Recall that an integral multicurve is a finite set of oriented curves with integral weights. By convention, a negative weight corresponds to reversing the orientation.




Figure 1. Proof of Lemma 8.8.

Lemma 8.9 *Suppose T is a geodesic triangulation of an orientable surface R in Ω_g . We orient each edge of T so that its x component is positive (for a vertical edge the orientation is not defined). Suppose W is a subset of the edges of T . Then there exists an integral multicurve Δ on R such that*

- (a) Δ is disjoint from the vertices of T and is transverse to the edges.
- (b) Δ crosses each edge of W at least once.
- (c) Each time Δ crosses an edge of $e \in W$, the crossing is from left to right (with respect to the orientation on the edge). If e is vertical, then all crossings must be from the same side.
- (d) For each edge e of T , the intersection number $i(\Delta, e) \leq n$, where n depends only on the genus g .

Proof. Each triangulated surface which has vertical edges is the limit of triangulated surfaces which do not. Hence, without loss of generality, we may assume that T has no vertical edges.

We define a directed graph $\mathcal{G}$ as follows (See Figure 2). The vertices of $\mathcal{G}$ are the connected components of $R \setminus W$. Two vertices A and B of $\mathcal{G}$ are connected by a directed edge of $\mathcal{G}$ if and only if there exists a saddle connection $\gamma \in W$ such that A is incident to γ from the left and B is incident to γ from the right.

We now claim that for each pair A, B of vertices of $\mathcal{G}$ there exists at least one directed path from A to B . Suppose this is not true. Let $\mathcal{A}$ be the set of vertices of $\mathcal{G}$ which can be reached from A by a directed path. Then $A \in \mathcal{A}$ and $B \in \mathcal{A}^c$; in particular $\mathcal{A}$ and $\mathcal{A}^c$ are both non-empty. Now let D be the closure of the connected components in $R \setminus W$ corresponding to vertices of $\mathcal{A}$. Then D is a subsurface of R , and ∂D consists of edges from W . For each edge $\gamma \in \partial D$ let $\epsilon_\gamma = +1$ if D is on the left of γ and -1 otherwise. Then, (since D is orientable) $\sum_{\gamma \in \partial D} \epsilon_\gamma \gamma = 0$. It follows that there exists $\gamma \in \partial D$ such that the x component of $\epsilon_\gamma \gamma$ is positive. Since the x component of γ is positive, this implies that $\epsilon_\gamma = 1$. Thus the directed edge of $\mathcal{G}$ corresponding to γ is directed from D to D^c . This means that some vertex of $\mathcal{G}$ outside of $\mathcal{A}$ can be reached from a vertex in $\mathcal{A}$ by a directed path, contradicting the definition of $\mathcal{A}$.




Figure 2. Lemma 8.9. The surface consists of three flat tori glued to each other as shown. The set W consists of the three dotted lines separating the tori. (The two dotted lines on the opposite sides of the figure are identified). The multicurve (in this case closed curve) Δ is drawn. The black dots are vertices of the graph $\mathcal{G}$ (used in the proof).

Thus, every pair of vertices of $\mathcal{G}$ can be connected by a directed path. Since $\mathcal{G}$ is a finite graph, this implies every directed edge of $\mathcal{G}$ is contained in a directed cycle of $\mathcal{G}$. Therefore there exists a finite union Δ of directed cycles such that every edge in $\mathcal{G}$ is contained in Δ . The length of Δ is bounded in terms of the size of $\mathcal{G}$, i.e. in terms of the genus. We can thus realize Δ as a multicurve on R with properties (a)-(d). $\square$

Lemma 8.10 *There exists a number $\rho_1 > 0$ such that for any $q \in \pi^{-1}(\mathcal{K})$ there exists $q' \in \pi^{-1}(\mathcal{K}) \cap \mathcal{F}^{uu}(q)$ and a path $\gamma : [0, \rho_1] \rightarrow \mathcal{F}^{uu}(q)$ such that $\gamma(0) = q$, $\gamma(\rho_1) = q'$, and for $t \in [0, \rho_1]$,*

$$\ell(\gamma(t)) \geq \ell(\gamma(0))\sqrt{1+t^2}, \quad (30)$$

and also

$$d_H(q, q') \leq C \int_0^{\rho_1} |\log(\ell(q)\sqrt{1+t^2})|^{1/2} dt. \quad (31)$$

Proof. Let $\tilde{S} = (\tilde{M}, \tilde{\omega})$ denote the double cover corresponding to q . Consider the Delaunay triangulation of $\tilde{S}$. By Lemma 8.8, all saddle connections of length at most $\sqrt{2}\ell(\tilde{S})$ belong to the Delaunay triangulation. Let W denote this set of saddle connections. Let Δ be the multicurve obtained by applying Lemma 8.9 to $\tilde{S}$ and W .

Let τ be the involution corresponding to q . Let $\Delta' = \Delta - \tau(\Delta)$. Then Δ' also has properties (a)-(d) of Lemma 8.9. In addition, $\tau(\Delta') = -\Delta'$.

Let $\rho_1 \in (0, \sqrt{2})$ be a constant which will be chosen later (depending only on the genus). We now use Δ' to define a path $\gamma : [0, \rho_1] \rightarrow \mathcal{Q}^1\mathcal{T}_g$ contained in $\pi^{-1}(\mathcal{K})$ and staying on the same leaf of $\mathcal{F}^{ss}$. We will actually define the path $\tilde{\gamma}(t)$ where for each t , $\tilde{\gamma}(t)$ is the double cover of $\gamma(t)$. Let $\tilde{S}$ denote the double cover of q , and set $\tilde{\gamma}(0) = \tilde{S}$. For each $t \in [0, \rho_1]$, $\tilde{\gamma}(t)$ is built from the same triangles as $\tilde{S}$, but for each edge e , we add $I(e, \Delta')t\ell(q)$ to the x -component, where $I(\cdot, \cdot)$ denotes the algebraic intersection number (cf. [MasSm91, §6]).

Note that we do not assert that the Delaunay triangulation of $\tilde{\gamma}(t)$ is the same as that of $\tilde{S} = \tilde{\gamma}(0)$. However, because of the form of Δ' , the involution τ acts on $\tilde{\gamma}(t)$, and we can let $\gamma(t)$ be the quotient by τ .

We now claim that (30) holds. Indeed, if $e \in W$ is a saddle connection in $\tilde{\gamma}(0)$ with vector (x_0, y_0) , then $x_0 \geq 0$, and on $\tilde{\gamma}(t)$, e has length

$$\sqrt{y_0^2 + (x_0 + I(e, \Delta')\ell(q)t)^2} \geq \sqrt{\ell(q)^2 + \ell(q)^2 t^2} \geq \ell(q)\sqrt{1+t^2} \quad (32)$$

Suppose $t_1, t_2 \in [0, \rho_1]$ and η are such that η is a saddle connection on $\tilde{\gamma}(t)$ for $t_1 < t < t_2$. Let $\ell_t(\eta)$ denote the length of η in the flat metric on $\tilde{\gamma}(t)$. Then,

$$\ell_t(\eta) \geq \ell_{t_1}(\eta) - |I(\eta, \Delta)|\ell(q)(t - t_1). \quad (33)$$

Now suppose z and w be any two zeroes of q . Let $\lambda_t(z, w)$ denote the shortest path between z and w on $\tilde{\gamma}(t)$, and let $|\lambda_t(z, w)|$ denote the length of $\lambda_t(z, w)$, i.e. the distance between z and w in the flat metric on $\tilde{\gamma}(t)$. Suppose $\lambda_0(z, w)$ is not an edge in W . Then, either $\lambda_0(z, w)$ is a saddle connection not in W , in which case $|\lambda_0(z, w)| \geq \sqrt{2}\ell(q)$, or $\lambda_0(z, w)$ is a union of at least two saddle connections, so that $|\lambda_0(z, w)| \geq 2\ell(q) \geq \sqrt{2}\ell(q)$. It now follows from (33) that for all $t \in [0, \rho_1]$,

$$|\lambda_t(z, w)| \geq (\sqrt{2} - nmt)\ell(q),$$

where n is the maximum intersection number of Δ' with a saddle connection in the Delaunay triangulation of $\tilde{S}$, and m is the maximal number of saddle connections in $\lambda_t(z, w)$. Note that both n and m are bounded by the genus. Now we choose ρ_1 so that $\sqrt{2} - nm\rho_1 \geq (1 + \rho_1)^2$. Then, if $\lambda_0(z, w)$ is not a saddle connection in $U \cap W$, then for all $t \in [0, \rho_1]$,

$$|\lambda_t(z, w)| \geq \ell(\gamma(0))\sqrt{1+t^2} \quad (34)$$

Now (30) follows from (32) and (34).

We now estimate the Hodge length of the path $\gamma : [0, \rho_1] \rightarrow \mathcal{Q}^1\mathcal{T}_g$. By Lemma 8.1 and Lemma 8.2,

$$\ell_\sigma(\gamma(t)) \geq \frac{C_g}{|\log(\ell(q)\sqrt{1+t^2})|},$$

where $\ell_\sigma(S)$ is defined to be the infimum over all simple closed curves α of $\ell_\sigma(\alpha)$ (and σ is the hyperbolic metric in the conformal class of S). By construction, the intersection number of Δ' with a short basis (see §8.2) of $\gamma(t)$ is bounded depending only on the genus. Therefore, by (21) and (22),

$$\|\gamma'(t)\|' \leq C|\log(\ell(q)\sqrt{1+t^2})|^{1/2},$$

where C depends only on the genus. Thus, if $q_1 = \gamma(\rho_1)$, then

$$d_H(q, q_1) \leq \int_0^{\rho_1} \|\gamma'(t)\|' dt \leq \int_0^{\rho_1} C|\log(\ell(q)\sqrt{1+t^2})|^{1/2} dt.$$

Thus (31) holds. □

Lemma 8.11 *There exists $\epsilon_0 > 0$ (depending on $\mathcal{K}$) such that for all $\epsilon < \epsilon_0$ and for all $q \in \pi^{-1}(\mathcal{K})$ with $\ell(q) < \epsilon$, there exists q' on the same leaf of $\mathcal{F}^{uu}$ as q with $\ell(q') \geq \epsilon$, and $d_H(q, q') \leq k\epsilon |\log \epsilon|^{1/2}$, where k depends only on $\mathcal{K}$.*

Proof. We define a sequence q_n as follows: let $q_0 = q$. If q_n has been defined already, we apply Lemma 8.10 with $q = q_n$, and define q_{n+1} to be the point q' guaranteed by Lemma 8.10. We obtain a sequence q_n with

$$\ell(q_{n+1}) \geq \ell(q_n)(1 + \rho_1^2)^{1/2} = \rho_2 \ell(q_n),$$

where we let $\rho_2 = (1 + \rho_1^2)^{1/2}$. Define t_n inductively by $t_0 = 0$, $t_{n+1} = t_n + \rho_1 \ell(q_n)$, $n \geq 0$. Then,

$$t_n = \rho_1 \sum_{k=0}^{n-1} \ell(q_k) \leq \rho_1 \sum_{k=0}^{n-1} \frac{\ell(q_n)}{\rho_2^{n-k}} = \rho_1 \ell(q_n) \frac{\rho_2^{-1} - \rho_2^{-n-1}}{1 - \rho_2^{-1}} \leq \frac{\rho_1 \ell(q_n)}{\rho_2 - 1}$$

Then,

$$t_{n+1} = t_n + \rho_1 \ell(q_n) \leq \frac{\rho_1 \rho_2}{\rho_2 - 1} \ell(q_n)$$

Thus, for $t \in [t_n, t_{n+1}]$,

$$\ell(q_n) \geq \frac{\rho_2 - 1}{\rho_1 \rho_2} t \equiv \rho_3 t$$

We have

$$d_H(q_n, q_{n+1}) \leq \int_{t_n}^{t_{n+1}} C |\log(\ell(q_n) \sqrt{1+t^2})|^{1/2} dt \leq \int_{t_n}^{t_{n+1}} C |\log(\rho_3 t \sqrt{1+t^2})|^{1/2} dt$$

Thus,

$$d_H(q_0, q_n) \leq \int_0^{t_n} C |\log(\rho_3 t \sqrt{1+t^2})|^{1/2} dt$$

We choose n so that $\ell(q_n)$ is comparable to ϵ . Let $q' = \gamma(t_n)$. Then, $\ell(q') \geq t_n \approx \epsilon$. Now the modified Hodge length of the path $q_0, q_1, \dots, q_n$ is $O(\epsilon |\log \epsilon|^{1/2})$. $\square$

Proof of Theorem 8.7. Since $\mathcal{K}$ is compact, the intersection of the multiple zero locus with $\pi^{-1}(\mathcal{K})$ is (contained in) finite union of hyperplanes $H_1, \dots, H_n$. Each hyperplane H_j has complex codimension 1. We can choose $\epsilon_0 > 0$ such that any two H_j which do not intersect in $\pi^{-1}(\mathcal{K})$ are at least ϵ_0 apart. Let $\delta_0 > 0$ be a lower bound on the angle between any two H_j which do intersect in $\pi^{-1}(\mathcal{K})$. Clearly ϵ_0 and δ_0 depend only on $\mathcal{K}$.

Let Z be the locus where our quadratic differential has either a zero of order at least 3 or two zeroes each of order at least 2. Then there exists a constant k_0 (depending only on δ_0 and thus only on $\mathcal{K}$) such that for any $q \in \pi^{-1}(\mathcal{K})$ and any $\epsilon > 0$, if the ball $B_E(q, \epsilon)$ intersects at least two hyperplanes H_j then $d_E(q, Z) < k_0 \epsilon$.

Take two points $q_1, q_2 \in \pi^{-1}(\mathcal{K})$ on the same leaf of $\mathcal{F}^{uu}$, with $d(q_1, q_2) = \epsilon$. Choose $k_3 > k_0$ (thus k_3 depends only on $\mathcal{K}$). We may also assume that $k_3 \epsilon < \epsilon_0$. By Lemma 8.11 (with $k_3 \epsilon$ in place of ϵ), there exist q'_1, q'_2 so that for $i = 1, 2$, we have $d_H(q_i, q'_i) \leq k_1 \epsilon |\log \epsilon|^{1/2}$, $d_E(q_i, q'_i) \leq k_2 \epsilon$, and $\ell(q_i) = k_3 \epsilon$, where k_1, k_2 depend only on $\mathcal{K}$. Since $k_3 > k_0$ and $k_3 \epsilon < \epsilon_0$, there exist ϵ_1, ϵ'_1 such that $k_3 \epsilon / 2 > \epsilon_1 > \epsilon'_1 > \epsilon / (2k_3)$, and for all $q \in \pi^{-1}(\mathcal{K})$, either $d_E(q, Z) < \epsilon_1$ or $B_E(q, \epsilon'_1)$ contains at most one hyperplane from the multiple zero locus.

Note that the intersection of Z with $\pi^{-1}(\mathcal{K})$ has complex codimension at least 2. Hence, the intersection of the ϵ_1 -neighborhood of Z with $\mathcal{F}^{uu}(q_1) \cap \pi^{-1}(\mathcal{K})$ is contained in the ϵ_1 -neighborhood of a finite union of hyperplanes, each of real codimension at least 2. Then there exists a constant k_4 depending only on $\mathcal{K}$, and a path γ connecting q'_1 to q'_2 of length at most $k_4\epsilon$, which avoids the ϵ_1 neighborhood of Z .

Now let $p_0 = q'_1$, and mark points p_i along γ which are $\epsilon'_1/2$ apart in the Euclidean metric. We have $p_n = q'_2$. Let B_i be the ball of Euclidean diameter $\epsilon'_1/2$ which contains p_i and p_{i+1} on its boundary. By construction B_i contains at most one hyperplane (which we will denote L) from the multiple zero locus. Note that by Lemma 8.1, Lemma 8.2 and (24), for any $p \in B_i$ and any tangent vector v at p , the modified Hodge norm of v can be estimated as

$$\|v\|'_H \leq C |\log d_E(p, L)|^{1/2} \|v\|_E, \quad (35)$$

where $\|v\|_E$ is the Euclidean norm of v , and $d(p, L)$ denotes the Euclidean distance between the point p and the hyperplane L .

Let p'_i be the farthest point in B_i from the hyperplane. Then, after connecting p_i and p'_i by a straight line path and using (35), we see that $d_H(p_i, p'_i) = O(\epsilon |\log \epsilon|^{1/2})$, and also $d_H(p'_i, p_{i+1}) = O(\epsilon |\log \epsilon|^{1/2})$. Thus, since the number of B_i along the path is bounded by a constant depending only on $\mathcal{K}$, we finally obtain

$$d_H(q'_1, q'_2) = O(d_E(q'_1, q'_2) |\log \epsilon|^{1/2})$$

□

8.5 The Non-expansion and Decay of the Hodge Distance

Theorem 8.12 *Suppose $q \in \mathcal{Q}^1\mathcal{T}_g$ and $q' \in \mathcal{Q}^1\mathcal{T}_g$ are in the same leaf of $\mathcal{F}^{ss}$. Then*

- (a) *There exists a constant $c_H > 0$ such that for all $t \geq 0$,*

$$d_H(g_t q, g_t q') \leq c_H d_H(q, q').$$

- (b) *Suppose $\epsilon > 2\epsilon_1$, where ϵ_1 is as in the definition of short basis. Let $K = \{q \in \mathcal{Q}^1\mathcal{T}_g : \ell(q) > \epsilon\}$. Suppose $d_H(q, q') < 1$, and $t > 0$ is such that*

$$\{s \in [0, t] : g_s q \in \Gamma K\} \geq (1/2)t. \quad (36)$$

Then for all $0 < s < t$,

$$d_H(g_s q, g_s q') \leq C e^{-cs} d_H(q, q'),$$

where c and C depend only on g and ϵ .

Proof. The statement (a) follows immediately from Theorem 8.3. For the second statement, let $\gamma : [0, \rho] \rightarrow \mathcal{Q}^1\mathcal{T}_g$ be a modified Hodge length minimizing path connecting q to q' (and staying in the same leaf of $\mathcal{F}^{ss}$). We assume that γ is parametrized so that $d_H(\gamma(u), q) = u$. By assumption, $\rho < 1$.

Let $N = 4c_H/(c_1\epsilon\rho)$, where c_1 is as in Theorem 8.7. For $0 \leq j \leq N$, let $u_j = jc_1\epsilon/(4c_H)$. We now claim that for all j there exists t_j such that for $s > t_j$ such that $g_sq \in \Gamma K$, we have for all $0 \leq u < u_j$,

$$d_E(g_s\gamma(u), g_sq) < \epsilon \sum_{k=0}^j 2^{-k}, \quad (37)$$

and

$$d_H(g_s\gamma(u), g_sq) < Ce^{-cs}u, \quad (38)$$

where C and c depend only on g and ϵ . The equations (37) and (38) will be proved by induction on j . If $j = 0$ then there is nothing to prove. Now assume $j \geq 1$, and (37) and (38) are true for all $u \leq u_{j-1}$. Suppose $u < u_j$. By (a), we have for all $t > 0$,

$$d_H(g_t\gamma(u), g_t\gamma(u_{j-1})) < c_H(u - u_{j-1}) < c_1\epsilon/4.$$

Therefore by Theorem 8.7, $d_E(g_t\gamma(u), g_t\gamma(u_{j-1})) < \epsilon/2$. Therefore, by the inductive assumption (37), for the $s > t_{j-1}$ such that $g_sq \in \Gamma K$, for all $u_{j-1} < u < u_j$, we have $\ell(g_s\gamma(u)) > \epsilon/2$. Thus, by (26), for all $u_{j-1} < u < u_j$, the modified Hodge norm of $\gamma'(u)$ is within a constant of the Hodge norm of $\gamma'(u)$. Then by Theorem 4.3, for all $u_{j-1} < u < u_j$,

$$d_H(g_t\gamma(u), g_t\gamma(u_{j-1})) < Ce^{-ct}d_H(\gamma(u), \gamma(u_{j-1})). \quad (39)$$

Therefore, by Theorem 8.7, there exists $t_j > 0$ (depending only on g and ϵ) such that for $s > t_j$ with $g_sq \in \Gamma K$, and $u_{j-1} \leq u < u_j$, (37) holds. Also (38) follows from (39). The induction terminates after finitely many steps (depending only on g and ϵ). Thus (b) holds. $\square$

9 The multiple zero locus

In this section we prove Theorem 6.1 and Theorem 6.2.

Let $\mathcal{K}$ be a compact subset of $\mathcal{T}_g$. We assume that $\mathcal{K}$ is contained in one fundamental domain for the action of Γ on $\mathcal{T}_g$ (and thus we can identify $\mathcal{K}$ with a subset of $\mathcal{M}_g = \mathcal{M}_g$). All of our implied constants will depend on $\mathcal{K}$.

Notation. Suppose $W \subset \mathcal{Q}^1\mathcal{T}_g$ and $s > 0$. Let $W(s)$ denote the set of $q \in \mathcal{Q}^1\mathcal{T}_g$ such that there exists $q' \in W$ on the same leaf of $\mathcal{F}^{uu}$ as q such that $d_H(q, q') < s$. Recall that for a subset $A \subset \mathcal{T}_g$, $Nbhd_r(A)$ denotes the set of points within Teichmüller distance r of A .

Lemma 9.1 *Suppose $U \subset \pi^{-1}(\mathcal{K})$, $\delta > 0$ and $t > 0$. Let $W = g_tU \cap \Gamma\pi^{-1}(\mathcal{K})$. Then,*

$$\mathbf{m}(Nbhd_2(\pi(W))) \leq C(\delta)\overline{\nu}(\eta^+(W(\delta))), \quad (40)$$

(By our convention, the constant $C(\delta)$ depends on $\mathcal{K}$ as well). Also

$$\overline{\nu}(\eta^+(W(1))) \leq C(\delta)\overline{\nu}(\eta^+(W(\delta))). \quad (41)$$

Proof. Suppose $x \in \text{Nbhd}_2(\pi(W))$. Then there exist $q_1 \in \pi^{-1}(x)$, $q_3 \in W$ and q_2 on the same leaf of $\mathcal{F}^u$ as q_3 and the same leaf of $\mathcal{F}^{ss}$ as q_1 with $d_{\mathcal{T}}(\pi(q_1), \pi(q_2)) = O(1)$, $d_{\mathcal{T}}(\pi(q_2), \pi(q_3)) = O(1)$. Then by Theorem 8.7, $d_H(q_2, q_3) \leq s$, where $s = O(1)$, so $q_2 \in W(s)$. Thus, every point of $\text{Nbhd}_2(\pi(W))$ is $O(1)$ distance along $\mathcal{F}^{ss}$ from a point of $W(s)$. Thus,

$$\mathbf{m}(\text{Nbhd}_2(\pi(W))) \leq C\overline{\nu}(\eta^+(W(s))).$$

Finally, $\overline{\nu}(\eta^+(W(s))) \leq C(\delta, s)\overline{\nu}(\eta^+(W(\delta)))$, (e.g. since d_H is equivalent to d_E). $\square$

The sets K_i and U_i . Let $K_1 \subset \mathcal{Q}^1\mathcal{M}_g$ be a compact set. (In our application, K_1 will be chosen disjoint from the multiple zero locus). Let $K_3 \subset K_2 \subset K_1$ and $1 \geq \delta > 0$ be such that if $q \in K_i$ and $d_H(q, q') < c_H\delta$ then $q' \in K_{i-1}$, where c_H is as in Theorem 8.12 (a). We assume $\overline{\mu}(K_3) > (1/2)$, where $\overline{\mu}$ is the normalized Lebesgue measure on $\mathcal{Q}^1\mathcal{M}_g$. For $T_0 > 0$ let $U_i = U_i(T_0) \subset \mathcal{Q}^1\mathcal{M}_g$ be such that for all $T > T_0$ and all $q \notin U_i$,

$$|\{t \in [0, T] : p(g_t q) \in K_i^c\}| < (1/2)T. \quad (42)$$

(Recall that p is the natural map from $\mathcal{Q}^1\mathcal{T}_g$ to $\mathcal{Q}^1\mathcal{M}_g$.) From the definition, we have $U_1 \subset U_2 \subset U_3$. By the ergodic theorem, for every $\theta > 0$ there exists $T_0 > 0$ such that $\overline{\mu}(U_3) < \theta$.

Let $\mathcal{K}_1 = \mathcal{K}$. We can choose compact subsets $\mathcal{K}_0, \mathcal{K}_1, \mathcal{K}_2$ of $\mathcal{T}_g$ such that for $0 \leq i \leq 3$, any $q \in \pi^{-1}(\mathcal{K}_i)$ and any q' on the same leaf of $\mathcal{F}^{uu}$ as q with $d_H(q, q') < c_H$, we have $q' \in \pi^{-1}(\mathcal{K}_{i+1})$. By (the lower bound in) Theorem 8.7 and Lemma 8.6, each $\mathcal{K}_i$ is compact. Let $U'_i = U_i \cap \pi^{-1}(\mathcal{K}_i)$. Note that $U'_1 \subset U'_2 \subset U'_3$.

Lemma 9.2 *Let U'_i , $1 \leq i \leq 3$ be as in the above paragraph. Then, for all $t > 0$,*

$$\mathbf{m}(\text{Nbhd}_2(\pi(g_t U'_1) \cap \Gamma\mathcal{K})) \leq C(\delta)e^{ht}\overline{\nu}(\eta^+(U'_2)),$$

and if $W(1)$ is defined as in Lemma 9.1 with $W = g_t U'_1 \cap \pi^{-1}(\Gamma\mathcal{K})$, then

$$\overline{\nu}(\eta^+(W(1))) \leq C'(\delta)e^{ht}\overline{\nu}(\eta^+(U'_2)).$$

In particular (c.f. Theorem 3.1), for any $\epsilon > 0$ it is possible to choose T_0 such that if U_1 is defined by (42) and $U'_1 = U_1 \cap \pi^{-1}(\mathcal{K}_1)$ then for all $t > T_0$,

$$\mathbf{m}(\text{Nbhd}_2(\pi(g_t U'_1) \cap \Gamma\mathcal{K})) \leq \epsilon e^{ht}, \quad (43)$$

and for $W = g_t U'_1 \cap \pi^{-1}(\Gamma\mathcal{K})$ we have

$$\overline{\nu}(\eta^+(W(1))) \leq \epsilon e^{ht}. \quad (44)$$

Proof. We will apply Lemma 9.1 to the set $W = g_t U'_1 \cap \pi^{-1}(\Gamma\mathcal{K})$. We claim that $W(\delta) \subset g_t U'_2$. Indeed, suppose $g_t q' \in W(\delta)$. Then there exists $g_t q \in W$ with

$$d_H(g_t q, g_t q') < \delta.$$

Then, by Theorem 8.12 (a), for all $0 \leq s \leq t$,

$$d_H(g_s q, g_s q') < c_H \delta \quad (45)$$

Since $W = g_t U'_1$, $g_t q \in W$ implies $q \in U'_1$. Then, assuming $t > T_0$, for at least half the values of $s \in [0, t]$,

$$g_s q \notin K_1. \quad (46)$$

Then, (45) and the definition of K_2 imply that for the $s \in [0, t]$ for which (46) holds, $g_s q' \notin K_2$. This implies $q' \in U_2$. Since $d_H(q, q') < c_H \delta < c_H$ and $q \in \pi^{-1}(\mathcal{K}) = \pi^{-1}(\mathcal{K}_1)$, we have $q' \in \pi^{-1}(\mathcal{K}_2)$. Thus, $q' \in U_2 \cap \pi^{-1}(\mathcal{K}_2) = U'_2$, and so $g_t q' \in g_t U'_2$. This implies the claim, and thus the first two statements of the Lemma.

The same argument as the proof of the claim shows that if $q \in U'_2$ and $q' \in \mathcal{F}^{ss}(q) \cap \pi^{-1}(\mathcal{K})$ with $d_H(q, q') < \delta$ then $q' \in U'_3$. This (together with Theorem 8.7) implies that there exists $C_1(\delta)$ such that

$$\overline{\nu}(\eta^+(U'_2)) \leq C_1(\delta) \overline{\mu}(U'_3).$$

Hence if we choose $T_0 > 0$ so that $\overline{\mu}(U_3) < C(\delta)C_1(\delta)\epsilon$, then (43) follows. Similarly, if we choose $T_0 > 0$ so that in addition $\overline{\mu}(U_3) < C'(\delta)C_1(\delta)\epsilon$, then (44) follows. $\square$

Proof of Theorem 6.2. Let $K' = K_1$, and let T_0 , U_1 and U'_1 be as in Lemma 9.2. Then, for $R > T_0$, and $x \in \mathcal{K}$,

$$B_R(x, \mathcal{K}, K') \subset \bigcup_{0 \leq t \leq R} \pi(g_t U'_1) \cap \Gamma \mathcal{K} = \bigcup_{n=1}^{R-1} \bigcup_{t \in [n, n+1]} \pi(g_t U'_1) \cap \Gamma \mathcal{K}.$$

Then,

$$\mathbf{m}(\text{Nbhd}_1(B_R(x, \mathcal{K}, K')))) \leq \sum_{n=0}^{R-1} \mathbf{m}(\text{Nbhd}_2(\pi(g_n U'_1) \cap \Gamma \mathcal{K})) \leq C\epsilon \sum_{n=0}^{R-1} e^{hnR},$$

where we have used (43). Since ϵ is arbitrary, Theorem 6.2 follows. $\square$

Lemma 9.3 Suppose $V \subset \mathcal{K}$ and $\delta' > 0$. Then, for t sufficiently large (depending on $\mathcal{K}$, V and δ'),

$$\mathbf{m}(\text{Nbhd}_1(\pi(g_t V) \cap \Gamma \mathcal{K}_0)) \leq C \overline{\nu}(\eta^+(V(\delta'))e^{ht}),$$

where C depends only on $\mathcal{K}$.

Proof. Let $Y = g_t V \cap \Gamma \mathcal{K}_0$. Choose T_0 so that (44) holds for $t > T_0$ and $\overline{\nu}(\eta^+(V(\delta')))$ instead of ϵ . Let U_1 , U'_1 and $W = g_t U'_1 \cap \pi^{-1}(\mathcal{K})$ be as in Lemma 9.2, so in particular, for $t > T_0$,

$$\overline{\nu}(\eta^+(W(1))) \leq \overline{\nu}(\eta^+(V(\delta'))e^{ht}). \quad (47)$$

We claim that there exist $T_1 > T_0$, depending only on $\mathcal{K}$ such that for $t > T_1$,

$$Y(1) = W(1) \cup g_t(V(\delta')). \quad (48)$$

Indeed, if $g_t q \in Y(1)$ then by definition there exists $q' \in V \subset \mathcal{K}$ with $d_H(g_t q, g_t q') < 1$, and q' is on the same leaf of $\mathcal{F}^{uu}$ as q . We consider two cases: either $q' \in U_1$ or $q' \notin U_1$. If $q' \in U_1$ then $q \in U'_1 = U_1 \cap \mathcal{K}$, hence $g_t q' \in g_t U'_1 \cap \Gamma \mathcal{K} = W$. Hence in this case, $g_t q \in W(1)$. If $q' \notin U_1$, then by (42) and Theorem 8.12 (b),

$$d_H(q, q') = d_H(g_{-t} g_t q, g_{-t} g_t q') \leq C e^{-ct} d_H(g_t q, g_t q') \leq C e^{-ct}.$$

We choose $T_1 > T_0$ so that $Ce^{-cT_1} < \delta'$. Thus, in this case, for $t > T_1$, $q \in V(\delta')$ and hence (48) follows.

Now, for $t > T_1$,

$$\overline{\nu}(\eta^+(Y(1))) \leq \overline{\nu}(\eta^+(W(1))) + \overline{\nu}(\eta^+(g_t(V(\delta')))) \leq 2\overline{\nu}(\eta^+(V(\delta')))e^{ht}$$

where we have used (47). We now apply Lemma 9.1 with $\delta = 1$. The lemma follows from (40). $\square$

Proof of Theorem 6.1. This follows immediately from Lemma 9.3, and the fact that we can choose an relatively open $V \subset \mathcal{K}$ and $\delta' > 0$ such that V contains the intersection of $\mathcal{K}$ with the multiple zero locus, and $\overline{\nu}(\eta^+(V(\delta')))$ is arbitrarily small, (see Theorem 3.1).

10 Volume Asymptotics

In this section we prove Theorem 1.2. Let $F_R(x, y)$ be as in §7. Fix $\epsilon_0 > 0$. We need the following

Theorem 10.1 *Given $x \in \mathcal{T}_g$ there exists $C = C(x)$ and $R_0 > 0$ such that for all $R > R_0$ and all $y \in \mathcal{T}_g$,*

$$F_R(x, y) \leq C(x)e^{hR}.$$

Proof of Theorem 1.2 assuming Theorem 10.1. By definition of F_R ,

$$\mathbf{m}(B_R(x)) = \int_{\mathcal{M}_g} F_R(x, y) d\mathbf{m}(y)$$

We multiply both sides by e^{-hR} and take the limit at $R \rightarrow \infty$. By Theorem 10.1 and we can apply the bounded convergence theorem to take the limit inside the integral. Now the theorem follows from Theorem 1.1. $\square$

A similar argument yields the following:

Theorem 10.2 *Suppose $x \in \mathcal{T}_g$, and $\mathcal{U} \subset S(x)$. Then, as $R \rightarrow \infty$,*

$$\mathbf{m}(B_R(x) \cap \text{Sect}_{\mathcal{U}}(x)) \sim \frac{1}{h} e^{hR} \Lambda(y) \int_{\mathcal{U}} \lambda^-(q) ds_x(q).$$

In the rest of this section we prove Theorem 10.1. Along the way, we prove Theorem 3.3.

Notation.

- Let $\mathcal{S}_g$ be the set of isotopy classes of integral *multicurves* on a surface of genus g .
- Given $x, y \in \mathcal{T}_g$ define

$$M_R(x, y) = \{\gamma \cdot y, \gamma \in \Gamma \mid \gamma \cdot y \in B_R(x)\},$$

so that $F_R(x, y) = \#(M_R(x, y))$.

The following result [Ker80] relates the ratios of extremal lengths to the Teichmüller distance:

Theorem 10.3 (Kerckhoff) *Given $x, y \in \mathcal{T}_g$, the Teichmüller distance between x and y is given by*

$$d_T(x, y) = \sup_{\beta \in \mathcal{S}_g} \log \left(\frac{\sqrt{\text{Ext}_\beta(x)}}{\sqrt{\text{Ext}_\beta(y)}} \right).$$

Estimating extremal length. We choose a constant C_g depending only on g such that for any surface $y \in \mathcal{T}_g$ there exists a pants decomposition $\mathcal{P} = \{\alpha_1, \dots, \alpha_{3g-3}\}$ of y such that the extremal lengths of the α_i are bounded from above by C_g .

Consider the Dehn-Thurston parameterization [HP92] of the set of multicurves

$$DT : \mathcal{S}_g \rightarrow (\mathbb{Z}_+ \times \mathbb{Z})^{3g-3}$$

defined by

$$DT(\beta) = (i(\beta, \alpha_i), \text{tw}(\beta, \alpha_i))_{i=1}^{3g-3},$$

where $i(\cdot, \cdot)$ denotes the geometric intersection number and $\text{tw}(\beta, \alpha_i)$ is the twisting parameter of β around α_i . See [HP92] for more details.

The following result is proved in [Mi96] (see Theorem 5.1, and equation (4.3)):

Theorem 10.4 (Minsky) *Suppose $y \in \mathcal{T}_g$, and let $\mathcal{P} = \mathcal{P}(y) = \{\alpha_1, \dots, \alpha_{3g-3}\}$ be any pants decomposition such that the extremal lengths of the α_i are bounded by C_g . Then given a simple closed curve β , $\text{Ext}_\beta(y)$ is bounded from above and below by*

$$\max_{1 \leq j \leq 3g-3} \left[\frac{i(\beta, \alpha_j)^2}{\text{Ext}_{\alpha_j}(y)} + \text{tw}^2(\beta, \alpha_j) \text{Ext}_{\alpha_j}(y) \right], \quad (49)$$

up to a multiplicative constant depending only on C_g and g .

Remark. The definition of the twist used in equation (4.3) in [Mi96] is different from the definition we are using here. We follow the definition used in [HP92]. Given a connected simple closed curve α on S_g , let $h_\alpha \in \Gamma$ denote the right Dehn twist around α . Then in terms of our notation,

$$\text{tw}(h_\alpha^r(\beta), \alpha) = \text{tw}(\beta, \alpha) + r \cdot i(\beta, \alpha). \quad (50)$$

Estimating the number of multicurves. Define

$$E(y, L) = \# \{ \alpha \in \mathcal{S}_g \mid \sqrt{\text{Ext}_\alpha(y)} \leq L \}$$

Let

$$G(y) = 1 + \prod_{\text{Ext}_\gamma(y) \leq \epsilon_0^2} \frac{1}{\sqrt{\text{Ext}_\gamma(y)}},$$

and the product ranges over all simple closed curves γ on the surface y with $\text{Ext}_\gamma(y) \leq \epsilon_0^2$.

Using Theorem 10.4, we obtain the following:

Theorem 10.5 *There exists a constant $C > 0$ such that for every $y \in \mathcal{T}_g$ and $L > 0$ we have*

$$E(y, L) \leq C \cdot G(y) \cdot L^{6g-6}.$$

Moreover, for any y , if $1/L$ is bounded by an absolute constant times the square root of the extremal length of the shortest curve on y , then

$$E(y, L) \leq CL^{6g-6}.$$

Sketch of the proof. In order to use the bound given by equation (49), first we fix a pants decomposition

$$\mathcal{P} = \{\alpha_1, \dots, \alpha_{3g-3}\}$$

on y such that $s_i = \sqrt{\text{Ext}_{\alpha_i}(y)} \leq C_g$ (depending only on g). In particular, this pants decomposition should include all small closed curves on y . Let $m_i = i(\beta, \alpha_i)$ and $t_i = \text{tw}(\beta, \alpha_i)$. So by equation (49), $\text{Ext}_\beta(y) \leq L^2$ implies that $\frac{m_i}{s_i} + |t_i|s_i = O(L)$. We use the following elementary lemma:

Lemma 10.6 *For $s > 0$, define $A_s(L)$ by*

$$A_s(L) = \{(a, b) \mid a, b \in \mathbb{Z}_+, a \cdot s + \frac{b}{s} \leq L\} \subset \mathbb{Z}_+ \times \mathbb{Z}_+.$$

Then for any $L > 0$, $|A_s(L)| \leq 4 \max\{s, 1/s\} \cdot L^2$. For $L > \max\{s, \frac{1}{s}\}$, we have $A_s(L) \leq 4L^2$.

Now applying the preceding lemma to the s_i , and using the Dehn-Thurston parameterization of multicurves, we get

$$E(y, L) \leq \prod_{i=1}^{3g-3} |A_{s_i}(L)| \leq C G(y) L^{6g-6}.$$

□

Proof of Theorem 3.3. This follows from the second part of Theorem 10.5. Recall that the extremal length can be extended continuously to a map

$$\text{Ext} : \mathcal{T}_g \times \mathcal{MF} \rightarrow \mathbb{R}_+$$

such that $\text{Ext}(x, t\lambda) = t^2 \text{Ext}(x, \lambda)$. Also the space $\mathcal{MF}$ has a piecewise linear integral structure, and elements of $\mathcal{S}_g$ are in one to one correspondence with the integral points. Hence,

$$\Lambda(x) = \text{Vol}\{\eta \in \mathcal{MF} \mid \sqrt{\text{Ext}_\eta(x)} \leq 1\} = \frac{\text{Vol}\{\eta \in \mathcal{MF} \mid \sqrt{\text{Ext}_\eta(x)} \leq L\}}{L^{6g-6}} = \lim_{L \rightarrow \infty} \frac{E(x, L)}{L^{6g-6}}, \quad (51)$$

where to justify the last equality use the fact that for any reasonably nice set A , as $L \rightarrow \infty$ the number of lattice points in the dilation LA is asymptotic to the volume of LA .

On the other hand, by the second assertion in Theorem 10.5 when L is big enough $E(y, L) \leq CL^{6g-6}$. Therefore, $\Lambda(y) \leq C$, where the bound does not depend on y . □

Notation. Fix a compact subset $K \subset \mathcal{M}_g$. For simplicity, we will also denote $\pi^{-1}(K) \subset \mathcal{T}_g$ by K . Note that for any $x \in K$ there exists a pants decomposition $\alpha = \{\alpha_i\}$ such that for any $1 \leq i \leq 3g-3$,

$$\frac{1}{C_K} \leq \sqrt{\text{Ext}_{\alpha_i}(x)} \leq C_K,$$

where C_K only depends on K . We call such a pants decomposition a *bounded* pants decomposition for $x \in K$.

We can show that for the elements in the thick part of the Teichmüller space, the Dehn-Thurston coordinates of a simple closed curve are bounded by its extremal length:

Lemma 10.7 *Let $\alpha = \{\alpha_1, \dots, \alpha_n\}$ be bounded pants decomposition of $x \in K$. Then there is a constant $c_1 > 0$, depending only on K , such that for any simple closed curve β on S_g ,*

$$\text{tw}(\beta, \alpha_i) \leq c_1 \cdot \sqrt{\text{Ext}_\beta(x)}, \quad i(\beta, \alpha_i) \leq c_1 \cdot \sqrt{\text{Ext}_\beta(x)}.$$

Proposition 10.8 *There exists a constant C_3 depending only on K such that for every $x \in K$, $y \in \mathcal{T}_g$ and $R > 0$ we have*

$$F_R(x, y) \leq C_3 E(y, e^R).$$

Our goal is to assign to any point $z = \gamma \cdot y \in M_R(x, y)$ a unique integral multicurve $\beta_z \in E(y, e^R)$. This would imply that

$$F_R(x, y) \leq C e^{(6g-6)R}.$$

(Note that we are assuming that x is in the compact part of $\mathcal{T}_g$ and $d(x, y) < R$; this implies that the shortest curve on y has extremal length at least constant times e^{-2R} , hence we may use the second statement of Theorem 10.5.)

The most natural candidate for β_z is $\gamma^{-1}\alpha$ where α is a fixed pants decomposition of x . However this correspondence is not one to one. Therefore, we need to modify the construction.

Given $z = \gamma \cdot y \in M_R(x, y)$, and $\alpha \in \mathcal{S}_g$ we have

$$\frac{\sqrt{\text{Ext}_\alpha(z)}}{\sqrt{\text{Ext}_\alpha(x)}} \leq e^R.$$

So by setting $L = e^R$, and $\beta = \gamma^{-1}(\alpha)$ we have

$$\sqrt{\text{Ext}_\beta(y)} \leq L \cdot \sqrt{\text{Ext}_\alpha(x)}. \quad (52)$$

Given $z = \gamma \cdot y \in M_R(x, y)$, let $\alpha(z)_i = \gamma^{-1}\alpha_i \in \mathcal{S}_g$. Then from (52) we get

$$\sqrt{\text{Ext}_{\alpha(z)_i}(y)} \leq C_K \cdot L.$$

Moreover, for $z_1 = \gamma_1 y, z_2 = \gamma_2 y \in \Gamma \cdot y$, we have $\alpha(z_1) = \alpha(z_2)$ if and only if there are $r_1, \dots, r_{3g-3} \in \mathbb{Z}$ such that

$$\gamma_1 = h_{\alpha_1}^{r_1} \cdots h_{\alpha_{3g-3}}^{r_{3g-3}} \cdot \gamma_2.$$

Lemma 10.9 *Let $\alpha = \{\alpha_1, \dots, \alpha_{3g-3}\}$ be a bounded pants decomposition on $x \in K$ and suppose $y_0 \in B_R(x)$. If $h_{\alpha_1}^{r_1} \cdots h_{\alpha_{3g-3}}^{r_{3g-3}}(y_0) \in B_R(x)$ then for $1 \leq i \leq 3g-3$*

$$|r_i| \cdot \sqrt{\text{Ext}_{\alpha_i}(y_0)} \leq C_2 e^R.$$

Here C_2 is a constant which only depends on K .

Sketch of the proof. Let $s_i = \sqrt{\text{Ext}_{\alpha_i}(y_0)}$, and $L = e^R$. The key point in the proof is Lemma 6.3 in [Mi96]. From this lemma, for any i , there exists a multicurve β_i of extremal length $b_i^2 = \text{Ext}_{\beta_i}(y_0)$ such that

$$i(\beta_i, \alpha_i) \geq cs_i \cdot b_i.$$

Since $d(x, y_0) \leq R$, $\sqrt{\text{Ext}_{\beta_i}(x)} \leq b_i \cdot L$. Also the assumption

$$d(h_{\alpha_1}^{-r_1} \cdots h_{\alpha_{3g-3}}^{-r_{3g-3}}(x), y_0) \leq R$$

implies that

$$\sqrt{\text{Ext}_{h_{\alpha_1}^{r_1} \cdots h_{\alpha_{3g-3}}^{r_{3g-3}}(\beta_i)}(x)} = \sqrt{\text{Ext}_{\beta_i}(h_{\alpha_1}^{-r_1} \cdots h_{\alpha_{3g-3}}^{-r_{3g-3}}(x))} \leq b_i \cdot L. \quad (53)$$

Then we have

1. Since $\sqrt{\text{Ext}_{\beta_i}(x)} \leq b_i \cdot L$, Lemma 10.7 implies that the twist and intersection coordinates of the curve β_i are bounded by a multiple of $b_i \cdot L$; in particular $|\text{tw}(\beta_i, \alpha_i)| = O(b_i \cdot L)$.
2. Similarly, by equation (53), applying Lemma 10.7 for $h_{\alpha_1}^{r_1} \cdots h_{\alpha_{3g-3}}^{r_{3g-3}}(\beta_i)$ on x with respect to $\{\alpha_i\}$ implies that

$$|\text{tw}(h_{\alpha_1}^{r_1} \cdots h_{\alpha_{3g-3}}^{r_{3g-3}}(\beta_i), \alpha_i)| = O(b_i \cdot L).$$

3. On the other hand, by the definition (see equation (50)) we have

$$\begin{aligned} |\text{tw}(h_{\alpha_1}^{r_1} \cdots h_{\alpha_{3g-3}}^{r_{3g-3}}(\beta_i), \alpha_i)| &= |r_i \cdot i(\alpha_i, \beta_i) + \text{tw}(\beta_i, \alpha_i)| \geq \\ &\geq |r_i| \cdot i(\alpha_i, \beta_i) - |\text{tw}(\beta_i, \alpha_i)|. \end{aligned}$$

So we have

$$|r_i| \cdot s_i \cdot b_i \leq \frac{1}{c} |r_i| \cdot i(\beta_i, \alpha_i) \leq C_2 L \cdot b_i \implies |r_i| \cdot s_i \leq C_2 L.$$

□

We remark that for any two disjoint simple closed curves η_1 and η_2 , and $m_1, m_2 > 0$, we have, by the definition of extremal length,

$$\sqrt{\text{Ext}_{m_1 \cdot \eta_1}(x)} + \sqrt{\text{Ext}_{m_2 \cdot \eta_2}(x)} \geq \sqrt{\text{Ext}_{m_1 \cdot \eta_1 + m_2 \cdot \eta_2}(x)}.$$

Corollary 1 *If the assumption of Lemma 10.9 holds, then for $\hat{\alpha} = \sum_{i=1}^{3g-3} |r_i| \cdot \alpha_i$ we have*

$$\sqrt{\text{Ext}_{\hat{\alpha}}(y_0)} \leq C_3 e^R.$$

Therefore, $\hat{\alpha}$ also defines a multicurve of extremal length bounded by e^{2R} . This completes the proof of Proposition 10.8, and thus in view of Theorem 10.5 the proof of Theorem 10.1.

A Appendix: Proof of Theorem 3.1.

In this section, $\mathcal{Q}^*\mathcal{T}_g$ denotes the space of non-zero quadratic differentials on marked compact surfaces of genus g , without restriction on area.

A local coordinate system near the multiple zero locus. Near the multiple zero locus, the coordinate system given by the period map (see Lemma 8.5) is singular. Instead we use an alternative coordinate system from [HuMas79].

Suppose $q_0 \in \mathcal{Q}^*\mathcal{T}_g$ has multiple zeroes, say $w_1, \dots, w_m$. Let m_i be the multiplicity of q_0 at w_i . Then, by [HuMas79, Proposition 3.1], for $1 \leq i \leq m$ there exists a choice of local coordinate z_i , mapping w_i to 0, such that any $q \in \mathcal{Q}^*\mathcal{T}_g$ near q_0 , on a neighborhood of w_i , has the form:

$$q = \left(z_i^{m_i} + \sum_{j=0}^{m_i-2} a_{ij} z_i^j \right) (dz_i)^2.$$

The coordinates z_i are uniquely determined by q . Fix some $\delta > 0$, and let w'_i be the point corresponding to $z_i = \delta$. For $q \in \mathcal{Q}^*\mathcal{T}_g$ near q_0 let $\tilde{M}$ denote the canonical double cover which makes the foliations corresponding to q orientable, and let σ be the involution of $\tilde{M}$ so that $\tilde{M}/\sigma$ is the surface M (with the flat structure given by q). Let $\Sigma_\delta = \sigma^{-1}(w'_1, \dots, w'_m)$, and let H denote the relative homology group $H_1(\tilde{M}, \Sigma_\delta, \mathbb{Z})$. Let H_{odd} denote the odd part of H under the action of the involution σ . Let $\gamma_1, \dots, \gamma_m$ be a integral basis for H_{odd} . Let

$$\lambda_i = \int_{\gamma_i} \sqrt{q}.$$

Suppose m_i is even. Let w_{ij} , $1 \leq j \leq m_i$ denote the zeroes of q which tend to w_i as $q \rightarrow q_0$. Let η_i be a fixed small circle in the coordinates z_i , centered at $z_i = 0$ (i.e. w_i). Then if q is sufficiently close to q_0 , η_i separates all the w_{ij} from the rest of the surface. Let $b_i = \int_{\eta_i} \sqrt{q}$. Then, a residue calculation shows that

$$b_i = a_{\frac{m_i}{2}-1} + \text{some polynomial in } a_{ij}, m_i - 2 \geq j \geq \frac{m_i}{2}. \quad (54)$$

Proposition A.1 (Hubbard-Masur) *If q_0 is not a square of an Abelian differential, then there exists a neighborhood $U \subset \mathcal{Q}^*\mathcal{T}_g$ of q_0 such that the a_{ij} and the λ_i are smooth local coordinates on U . If q_0 is the square of an Abelian differential, then there exists a neighborhood U of q_0 such that under the constraint $\sum_i b_i = 0$, the a_{ij} and the λ_i are local coordinates on U .*

In both cases, near q_0 , the natural projection map $\hat{\pi} : \mathcal{Q}^\mathcal{T}_g \rightarrow \mathcal{T}_g$ is a submersion in these coordinates.*

Proof. See [HuMas79, Proposition 4.7]. □

The constraint $\sum_i b_i = 0$ appears because in the case when q_0 is the square of an Abelian differential, the part of the surface which is outside all the circles η_i has an orientable foliation. In that case, in view of (54), we may drop $a_{\frac{m_i}{2}-1}$ for some i from the coordinate system.

We introduce a covering of $\mathcal{Q}^*\mathcal{T}_g$ by open sets U_α , such that each U_α is the neighborhood U of Proposition A.1, for some $q_0 \in \mathcal{Q}^*\mathcal{T}_g$. We may assume that the U_α are invariant under the operation of multiplying the quadratic differential by a real number. We also assume that covering

is uniformly locally finite on compact sets, i.e. that for any compact set $\mathcal{K} \in \mathcal{T}_g$, there exist a number N depending only on $\mathcal{K}$ such that each point in $\hat{\pi}^{-1}(\mathcal{K})$ belongs to at most N sets U_α .

Let ψ_α be a partition of unity subordinate to U_α . For $q, q' \in U_\alpha$, let $\{a_{ij}, \lambda_i\}$ be the coordinates of q , and $\{a'_{ij}, \lambda'_i\}$ be the coordinates of q' . Let $D_\alpha(q, q') = \sum_{ij} |a_{ij} - a'_{ij}| + \sum_i |\lambda_i - \lambda'_i|$. For q and q' sufficiently close so that they belong to the same U_α , let $D(q, q') = \sum_\alpha \psi_\alpha D_\alpha(q, q')$.

Recall $d_{\mathcal{T}}(x, y)$ denotes the Teichmüller distance between x and y , and $S(x)$ denotes the sphere of unit area quadratic differentials which are holomorphic at x .

Lemma A.2 *Suppose x, y in $\mathcal{T}_g$ are sufficiently close and belong to a compact set $\mathcal{K}$. Then, there exist $c > 1$ depending only on $\mathcal{K}$ such that*

$$c^{-1} \inf_{q \in S(x)} D(q, \hat{\pi}^{-1}(y)) \leq d_{\mathcal{T}}(x, y) \leq \sup_{q \in S(x)} D(q, \pi^{-1}(y)). \quad (55)$$

Proof. Let K be the pullback of $\mathcal{K}$ from $\mathcal{T}_g$ to $\mathcal{Q}^1\mathcal{T}_g$. Then K is also compact. Near any $q \in K$, by Proposition A.1, the map $p : \mathcal{Q}^*\mathcal{T}_g \rightarrow \mathcal{T}_g$ is a submersion, and therefore can be approximated by a linear map, for which (55) is immediate. The rest follows by compactness. $\square$

Let β_α be the measure on $U_\alpha \subset \mathcal{Q}^*\mathcal{T}_g$ given by $\prod_{i,j} da_{ij} \prod_i d\lambda_i$. Let $\beta = \sum_\alpha \psi_\alpha \beta_\alpha$.

For $V \subset \pi^{-1}(x)$, let $V^* \subset \mathcal{Q}^*\mathcal{T}_g$ denote the set of all quadratic differentials which have the same horizontal foliation as some $q \in V$.

Lemma A.3 *For every $\delta > 0$, any $x \in \mathcal{T}_g$ and any $q_0 \in \mathcal{Q}^*\mathcal{T}_g$ there exists an open subset V of $\pi^{-1}(x)$ containing q_0 , such that for all sufficiently small $\epsilon > 0$,*

$$\beta(V^* \cap \pi^{-1}(B(x, \epsilon))) < \delta \epsilon^{6g-6}.$$

Proof. Fix $\delta' > 0$. Recall that by a horosphere (e.g. leaf of $\mathcal{F}^{ss}$) in $\mathcal{Q}^*\mathcal{T}_g$ we mean the set of quadratic differentials with a fixed horizontal foliation. Since the foliation by horospheres is continuous, there exists a neighborhood V of $q_0 \in \pi^{-1}(x)$ such that $V \subset B_D(q_0, \delta')$, where B_D denotes a ball in the metric D . Then, for $\epsilon > 0$ sufficiently small, and in view of Lemma A.2, there exist constants C_1, C_2 such that

$$\pi^{-1}(V^* \cap B(x, \epsilon)) \subset \{q \in \mathcal{Q}^*\mathcal{T}_g : D(q, B_D(q_0, C_2\delta')) \cap \pi^{-1}(x)\} < C_1\epsilon. \quad (56)$$

Since $\pi^{-1}(x)$ is smooth, the β -measure of the set on the right of (56) is $\epsilon^{6g-6}c(\delta')$, where $c(\delta') \rightarrow 0$ as $\delta' \rightarrow 0$. This implies the lemma. $\square$

Proposition A.4 *The measure μ on $\mathcal{Q}^*\mathcal{T}_g$ is absolutely continuous with respect to β , and for any compact subset K of $\mathcal{T}_g$ there exists a constant C such that for all $q \in \pi^{-1}(K)$, $\left| \frac{d\mu}{d\beta}(q) \right| < C$. Also, away from the multiple zero locus, $\left| \frac{d\mu}{d\beta}(q) \right|$ is a smooth non-vanishing function of q .*

Remark. It is also possible to show that there exists a constant C' depending only on K such that for all $x \in \pi^{-1}(K)$, one has $C' < \left| \frac{d\mu}{d\beta}(x) \right|$. Since we do not need this, we will omit the proof.

Proof of Theorem 3.1, assuming Proposition A.4. By Lemma A.2, $\beta(\pi^{-1}(B(x, \epsilon))) = O(\epsilon^{6g-6})$. Therefore, by Proposition A.4, there exist $0 < c_1 < c_2$ such that

$$c_1 \epsilon^{6g-6} < \mu(\pi^{-1}(B(x, \epsilon))) < c_2 \epsilon^{6g-6}. \quad (57)$$

Now,

$$ds_x(V) = \lim_{\epsilon \rightarrow 0} \frac{\mu(V^* \cap \pi^{-1}(B(x, \epsilon)))}{\mu(\pi^{-1}(B(x, \epsilon)))} \leq \limsup_{\epsilon \rightarrow 0} \frac{C\beta(V^* \cap \pi^{-1}(B(x, \epsilon)))}{\mu(\pi^{-1}(B(x, \epsilon)))} \leq \frac{C\delta\epsilon^{6g-6}}{c_1\epsilon^{6g-6}}.$$

where for the first inequality we used Proposition A.4, and for the last estimate we used Lemma A.3 and (57). Since δ is arbitrary, the theorem follows. $\square$

The rest of the section will consist of the proof of Proposition A.4. To simplify notation, we work with one zero of q_0 at a time. We may write:

$$z^m + \sum_{i=0}^{m-1} a_i z^i = \prod_{j=1}^m (z - z_j).$$

Then the a_i are symmetric polynomials in the z_j . It is well known that the Jacobian of the map from the zeroes z_j 's to the coefficients a_i 's is the Vandermonde determinant. In other words, if we use the notation $\frac{\partial(y_1, \dots, y_n)}{\partial(x_1, \dots, x_n)}$ for the Jacobian determinant of the matrix $\left\{ \frac{\partial y_i}{\partial x_j} \right\}_{1 \leq i, j \leq n}$, we have:

$$\frac{\partial(a_0, \dots, a_{m-1})}{\partial(z_1, \dots, z_m)} = \prod_{i < j} (z_i - z_j). \quad (58)$$

(This follows from comparing degrees and noting the anti-symmetry of the determinant).

We may choose a basis for the homology relative to the zeroes. This amounts to choosing a spanning subtree T from the complete graph connecting the zeroes z_j . We may choose the tree in such a way that for any z_i and z_j , $|z_i - z_j|$ is within a multiplicative constant (depending only on m) of the length of the path in T connecting z_i and z_j . Let the (oriented) edges of T be $e_1, \dots, e_{m-1}$. Let e_k^+ be the zero at the head of e_k , and e_k^- the zero at the tail of e_k . We write $\vec{e}_k = e_k^+ - e_k^-$ (so $\vec{e}_k \in \mathbb{C}$).

In our setting, we need to restrict to $a_{m-1} = 0$. Note that $a_{m-1} = z_1 + \dots + z_m$. It follows that

$$\frac{\partial(a_0, \dots, a_{m-2})}{\partial(\vec{e}_1, \dots, \vec{e}_{m-1})} = \frac{\partial(a_0, \dots, a_{m-1})}{\partial(\vec{e}_1, \dots, \vec{e}_{m-1}, a_{m-1})} = \frac{\partial(a_0, \dots, a_{m-1})}{\partial(z_1, \dots, z_m)} \frac{\partial(z_1, \dots, z_m)}{\partial(\vec{e}_1, \dots, \vec{e}_{m-1}, a_{m-1})} = \prod_{i < j} (z_i - z_j), \quad (59)$$

where we have used (58).

If z_j is a zero, and e is an edge of T , let

$$d_+(z_j, e) = \max(|z_j - e^-|, |z_j - e^+|).$$

We will need the following combinatorial lemma:

Lemma A.5

$$\prod_{e \in T} \prod_{p=1}^m d_+(z_p, e)^{1/2} \leq C \prod_{i < j} |z_i - z_j|, \quad (60)$$

where C depends only on m .

Proof of Lemma A.5. Let e' be the longest edge of T . If we cut along e' , we separate the tree into two subtrees T_1 and T_2 say of size m_1 and m_2 . If $z_i \in T_1$ and $z_j \in T_2$, then by the assumption on T , $|z_i - z_j|$ is comparable to the length of the path in the tree connecting z_i to z_j . This path contains e' , which is by assumption the longest edge in the tree. Therefore, if $z_i \in T_1$ and $z_j \in T_2$ then $|z_i - z_j|$ is within a multiplicative constant of e' . Hence, the left hand side of (60) is within a multiplicative constant of

$$|e'|^{m_1 m_2} \prod_{\substack{i < j \\ z_i, z_j \in T_1}} |z_i - z_j| \prod_{\substack{i < j \\ z_i, z_j \in T_2}} |z_i - z_j|.$$

To estimate the right hand side of (60), note that each factor of the form $d_+(z_p, e)^{1/2}$ where $z_p \in T_1$ and either $e \in T_2$ or $e = e'$ is within a multiplicative constant of $|e'|^{1/2}$. The number of such factors is $m_1 m_2$. Also the same factors appear when $z_p \in T_2$ and $e \in T_1$ or $e = e'$. Then the right hand side of (60) is within a multiplicative constant of

$$|e'|^{m_1 m_2} \prod_{e \in T_1} \prod_{z_p \in T_1} d_+(z_p, e)^{1/2} \prod_{e \in T_2} \prod_{z_p \in T_2} d_+(z_p, e)^{1/2}$$

The estimate (60) now follows by induction. □

Lemma A.6 Suppose $z_1, \dots, z_m$ are in some bounded set K . Let

$$\Omega_k = \int_{e_k^-}^{e_k^+} \sqrt{(z - z_1) \dots (z - z_n)} dz,$$

so that Ω_k is the holonomy of the edge e_k . Then,

$$\left| \frac{\partial(\Omega_1, \dots, \Omega_{n-1})}{\partial(\vec{e}_1, \dots, \vec{e}_{m-1})} \right| \leq C \prod_{i < j} |z_i - z_j|,$$

where C depends only on m and K .

Proof. Suppose for the moment that $z_1, \dots, z_m$ are independent variables. We claim that

$$\left| \frac{\partial \Omega_k}{\partial z_j} \right| \leq C \prod_{p=1}^n d_+(z_p, e_k)^{1/2}, \quad (61)$$

where C depends only on m and K . Indeed,

$$\frac{\partial \Omega_k}{\partial z_j} = \int_{e_k^-}^{e_k^+} \frac{\sqrt{(z - z_1) \dots (z - z_n)}}{2(z - z_j)} dz$$

If we write the numerator in the integral as $P(z - z_j)^{1/2}$, then P is bounded by a constant times

$\prod_{\substack{p=1 \\ p \neq j}}^n d_+(z_p, e_k)^{1/2}$. Then, the integral is bounded by

$$\frac{|P|}{2} \left| \int_{e_k^-}^{e_k^+} \frac{dz}{(z - z_j)^{1/2}} \right|$$

If $d_+(z_j, e_k) > \frac{1}{2}|e_k^+ - e_k^-|$,

$$\left| \int_{e_k^-}^{e_k^+} \frac{dz}{(z - z_j)^{1/2}} \right| \leq C \frac{|e_k^+ - e_k^-|}{d_+(z_j, e_k)^{1/2}} \leq C' |e_k^+ - e_k^-|^{1/2}$$

(where we have used the assumption that $|e_k^+ - e_k^-|$ is bounded). If $d_+(z_j, e_k) < \frac{1}{2}|e_k^+ - e_k^-|$, then

$$\left| \int_{e_k^-}^{e_k^+} \frac{dz}{(z - z_j)^{1/2}} \right| \leq C'' |e_k^+ - e_k^-|^{1/2}.$$

where C'' depends only on m and K . Thus,

$$\left| \frac{\partial \Omega_k}{\partial z_j} \right| \leq C |e_k^+ - e_k^-|^{1/2} \prod_{p \neq j} d_+(z_p, e_k)^{1/2}.$$

But by the choice of the tree T , for any j , $d_+(z_j, e_k) \geq c|e_k^+ - e_k^-|$, where c depends only on m . Thus, (61) holds.

Since $\sum_{j=1}^m z_j = 0$, there are $m-1$ linearly independent z_j 's and therefore we can express $z_j = \sum_{k=1}^{m-1} \eta_{jk} \vec{e}_k$, where the η_{jk} are bounded depending only on m . Then, we get

$$\left| \frac{\partial \Omega_j}{\partial \vec{e}_k} \right| \leq C' \prod_{p=1}^m d_+(z_p, e_k)^{1/2}.$$

Lemma A.6 now follows from Lemma A.5. $\square$

Proof of Proposition A.4. It is enough to show that for any α , $|\frac{\partial \mu}{\partial \beta_\alpha}| \leq C$ on U_α . As above, we work with one zero of q_0 at a time. By (59) and Lemma A.6,

$$\left| \frac{\partial(\Omega_1, \dots, \Omega_{m-1})}{\partial(a_0, \dots, a_{m-2})} \right| = \left| \frac{\partial(\Omega_1, \dots, \Omega_{m-1})}{\partial(\vec{e}_1, \dots, \vec{e}_{m-1})} \right| \left| \frac{\partial(\vec{e}_1, \dots, \vec{e}_{m-1})}{\partial(a_0, \dots, a_{m-2})} \right| \leq C, \quad (62)$$

where C depends only on m and K .

Let $\mu' = d\Omega_1 d\bar{\Omega}_1 \dots d\Omega_{m-1} d\bar{\Omega}_{m-1}$, and let $\beta' = da_0 d\bar{a}_0 \dots da_{m-2} d\bar{a}_{m-2}$. Then, by (62),

$$\frac{d\mu'}{d\beta'} = \left| \frac{\partial(\Omega_1, \dots, \Omega_{m-1})}{\partial(a_0, \dots, a_{m-2})} \right|^2 \leq C^2.$$

Now,

$$\frac{d\mu}{d\beta_\alpha} = \prod_j \frac{d\mu'_j}{d\beta'_j},$$

where the product is over distinct zeroes of q_0 . Now Proposition A.4 follows from (62). $\square$

References

- [Ac60] Robert D. M. Accola. Differentials and Extremal Length on Riemann Surfaces. Proceedings of the National Academy of Sciences of the United States of America, Vol. 46, No. 4. (Apr. 15, 1960), pp. 540-543.
- [Ath06] J. Athreya. *Quantitative recurrence and large deviations for Teichmüller geodesic flow*. Geom. Dedicata 119 (2006), 121–140.
- [Bl61] C. Blatter. *Über Extremallängen auf geschlossenen Flächen*. (German) Comment. Math. Helv. 35 1961 153–168.
- [CSF80] I.P. Cornfeld, Ya. G. Sinai, S.V. Fomin, Ergodic theory, “Nauka”, Moscow, 1980.
- [EM01] A. Eskin and H. Masur. Asymptotic formulas on flat surfaces. *Ergodic Theory Dynam. Systems*, **21** (2001), 443–478.
- [EMc93] A. Eskin and C. McMullen, *Mixing, counting, and equidistribution in Lie groups*. Duke Mathematical Journal, 71 (1993), no. 1, 181–209.
- [Fa73] Fay J.D. *Theta functions on Riemann surfaces*. Lecture Notes in Mathematics no. 352, Springer 1973.
- [Fo02] G. Forni, *Deviation of ergodic averages for area-preserving flows on surfaces of higher genus*. Ann. of Math. (2), 155 (2002), no. 1, 1–103.
- [HP92] J. L. Harer and R. C. Penner. *Combinatorics of Train Tracks*, volume 125 of *Annals of Math. Studies*. Princeton University Press, 1992.
- [HuMas79] J. Hubbard, H. Masur, *Quadratic differentials and foliations*. Acta Math. 142 (1979), no. 3-4, 221–274.
- [Ka90a] V. Kaimanovich, “Bowen-Margulis and Patterson measures on negatively curved compact manifolds.” Dynamical systems and related topics (Nagoya, 1990), 223–232, Adv. Ser. Dynam. Systems, 9, World Sci. Publ., River Edge, NJ, 1991.
- [Ka90b] V. Kaimanovich, “Invariant measures of the geodesic flow and measures at infinity on negatively curved manifolds.” Hyperbolic behaviour of dynamical systems (Paris, 1990). Ann. Inst. H. Poincaré Phys. Théor. 53 (1990), no. 4, 361–393.
- [Ker80] S. Kerckhoff. The asymptotic geometry of Teichmüller space. *Topology* **19**(1980), 23–41.
- [Ko96] M. Kontsevich, Lyapunov exponents and Hodge theory, “Mathematical Beauty of Physics”, Saclay, 1996.
- [KZ03] M. Kontsevich, A. Zorich, Connected components of the moduli spaces of Abelian differentials with prescribed singularities, *Inventiones mathematicæ*, 153(2003), no.3, 631-678.
- [LM] E. Lindenstrauss, M. Mirzakhani. *Ergodic theory of the space of measured laminations*, Int. Math. Res. Not. IMRN (2008), no. 4, Art. ID rnm126, 49 pp.

- [Mar70] G.A. Margulis, On some aspects of the theory of Anosov flows, Ph.D. Thesis, 1970, Springer, 2003.
- [Mas76] H. Masur, *Extension of the Weil-Peterson metric to the boundary of Teichmüller space*. Duke Math. J. 43 (1976), no. 3, 623–635.
- [Mas82] H. Masur, *Interval exchange transformations and measured foliations*. Ann. of Math. (2) 115 (1982), no. 1, 169–200.
- [MasSm91] H. Masur and J. Smillie *Hausdorff Dimension of Sets of Nonergodic Measured Foliations* Ann. of Math., Second Series, Vol. 134, No. 3 (Nov., 1991), pp. 455–543
- [Mi93] Y. Minsky. Teichmüller geodesics and ends of hyperbolic 3-manifolds. . Topology, 32 (1993), 625–647.
- [Mi96] Y. Minsky. Extremal length estimates and product regions in Teichmüller space. Duke Math. J. 83 (1996), no. 2, 249–286.
- [Ra05] K. Rafi. *Thick-thin decomposition of quadratic differentials*. preprint, 2005.
- [Ra07] K. Rafi. *A combinatorial model for the Teichmüller metric*. Geom. Funct. Anal., 17(3), 936–959, 2007.
- [Si94] Ya. G. Sinai, Topics in ergodic theory, Princeton University Press, 1994.
- [Ve78] W. Veech, Interval exchange transformations. J. Analyse Math. 33 (1978), 222–272.
- [Ve82] W. Veech, Gauss measures for transformations on the space of interval exchange maps, Annals of Mathematics, 15(1982), 201–242.
- [Ve86] W. Veech, The Teichmüller geodesic flow, Annals of Mathematics, 1986.

Department of Mathematics, Yale University, New Haven CT, USA;
jathreya@gmail.com

Department of Mathematics, Rice University, MS 136, 6100 Main Street, Houston, Texas 77251-1892 and The Steklov Institute of Mathematics, Russian Academy of Sciences, Gubkina str. 8, 119991, Moscow, Russia;
aib1@rice.edu and bufetov@mi.ras.ru

Department of Mathematics, University of Chicago, Chicago IL 60637, USA;
eskin@math.uchicago.edu

Department of Mathematics, Stanford University, Stanford CA 94305 USA;
mmirzakh@math.stanford.edu